

Parallelism between Moore-Penrose inverse and Aluthge transformation of operators

*M. R. Jabbarzadeh, H. Emamalipour and M. Sohrabi Chegeni**

In this paper we study some parallelisms between \dagger -Aluthge transform and binormal operators on a Hilbert space via the Moore-Penrose inverse. Moreover, we give some applications of these results on the Lambert multiplication operators acting on $L^2(\Sigma)$.

1. INTRODUCTION AND PRELIMINARIES

Let $B(H)$ denote the C^* -algebra of all bounded linear operators on a complex Hilbert space H . We write $\mathcal{N}(T)$ and $\mathcal{R}(T)$ for the null-space and the range of an operator $T \in B(H)$, respectively. Recall that for $T \in B(H)$, there is a unique factorization $T = U|T|$, where $\mathcal{N}(T) = \mathcal{N}(U) = \mathcal{N}(|T|)$, U is a partial isometry, i.e. $UU^*U = U$, and $|T| = (T^*T)^{1/2}$ is a positive operator. This factorization is called the polar decomposition of T . Note that $T = |T^*|U = \sqrt{|T^*|}U\sqrt{|T|}$. More generally, $T = |T^*|^p U |T|^{1-p}$ for $p \in (0, 1)$; see e.g. [13, Theorem 2.7]. If $T = U|T|$ is the polar decomposition of $T \in B(\mathcal{H})$, then $\tilde{T} = |T|^{1/2}U|T|^{1/2}$ is called the Aluthge transformation of T . Let $\text{CR}(H)$ be the set of all bounded linear operators on H with closed range. For $T \in \text{CR}(H)$, the Moore-Penrose inverse of T , denoted by T^\dagger , is the unique operator T^\dagger that satisfies the following:

$$TT^\dagger T = T, \quad T^\dagger TT^\dagger = T^\dagger, \quad (TT^\dagger)^* = TT^\dagger, \quad (T^\dagger T)^* = T^\dagger T.$$

*Corresponding author. Morteza Sohrabi Chegeni
2010 Mathematics Subject Classification. Primary: 47B20; Secondary: 47B38.
Keywords and Phrases. Aluthge transformation, Moore-Penrose inverse, polar decomposition, conditional expectation.

We recall that T^\dagger exists if and only if $T \in \text{CR}(H)$. Note that if $T \in \text{CR}(H)$, then T^* , $|T|$ and T^\dagger have closed range. If $T = U|T|$ is invertible, then $T^{-1} = T^\dagger$, U is unitary and so $|T|$ is invertible. It is a classical fact that the polar decomposition of T^* is $U^*|T^*|$. It is easy to check that $U^*|T^*|^\dagger$ and $|T^\dagger|^\frac{1}{2}U^*|T^\dagger|^\frac{1}{2}$ are the polar decomposition and Aluthge transformation of T^\dagger , respectively. For other important properties of T^\dagger see [2, 10, 17].

An operator $T \in B(H)$ is said to be binormal if $[|T|, |T^*|] = 0$, where $[A, B] = AB - BA$ for operators A and B . The numerical range $W(T)$ of an operator $T \in B(H)$ is defined by $W(T) = \{\langle Tx, x \rangle : \|x\| = 1\}$. Also, $\omega(T) = \sup\{|\lambda| : \lambda \in W(T)\}$ and $Sp(T)$ denote the numerical radius and spectrum of T , respectively.

Study of Moore-Penrose inverse and Aluthge transformation of bounded linear operators has a long history. In this paper, we introduce \dagger -Aluthge transformation which is parallel to Aluthge transformation. Then we investigate some connections and parallelisms between \dagger -Aluthge transformation and binormal operators via the Moore-Penrose inverse. In section 2, firstly, we give a necessary and sufficient condition to the quasinormality of T^\dagger . We show that if T is onto, then T^* is quasinormal if and only if T^\dagger is quasinormal. Afterward, we give a formula for $(\widetilde{T^*})^\dagger$ when T is binormal. Also, we prove that T^* is quasinormal if and only if $(T^\dagger)^* = (\widetilde{T^*})^\dagger$, whenever T is onto. Moreover, we briefly discuss some classical results on the spectrum, numerical range and numerical radius via the Moore-Penrose inverse and Aluthge transformation. In section 3, we obtain some applications of these results to the Lambert multiplication operator M_wEM_u on $L^2(\Sigma)$, where E is the conditional expectation operator with respect to a sub-sigma algebra $\mathcal{A} \subseteq \Sigma$. In addition, we determine lower and upper bounds estimates for the numerical range of $(M_wEM_u)^\dagger$.

2. ON SOME CHARACTERIZATIONS OF T^\dagger

For any closed subspace M of H , let P_M denote the orthogonal projection onto M . For $T \in \text{CR}(H)$, we shall make use of the following general properties of T^* , \widetilde{T} , T^\dagger and their polar decompositions. For proofs and discussions of these facts see [10, 9, 12, 20, 22].

- P(1) $\widetilde{T^\dagger} = |T^\dagger|^\frac{1}{2}U^*|T^\dagger|^\frac{1}{2}$;
- P(2) For $\lambda > 0$, $\lambda \in Sp(T)$ if and only if $\lambda^{-1} \in Sp(T^\dagger)$;
- P(3) $|T^\dagger| = |T^*|^\dagger$ and $|T^\dagger|^\frac{1}{2} = (|T^*|^\frac{1}{2})^\dagger$;
- P(4) $|T^*|^\frac{1}{2}(|T^*|^\frac{1}{2})^\dagger = P_{R(|T^*|)} = (|T^*|^\frac{1}{2})^\dagger|T^*|^\frac{1}{2}$;
- P(5) If T is binormal, then $P_{R(|T^*|)}P_{R(|T|)} = P_{R(|T|)}P_{R(|T^*|)}$;

- P(6) $U^*P_{R(|T^*|)} = U^* = P_{R(|T|)}U^*$;
P(7) $U^*U = P_{R(|T|)}$ and $UU^* = P_{R(|T^*|)}$;
P(8) $|T^\dagger|^{\frac{1}{2}}P_{R(|T^*|)} = |T^\dagger|^{\frac{1}{2}}$;
P(9) $U^*(|T^*|^\dagger)^{\frac{1}{2}} = (|T|^\dagger)^{\frac{1}{2}}U^*$;
P(10) $UU^*|T^*|^\dagger = |T^*|^\dagger$;
P(11) $(T^\dagger)^* = |T^*|^\dagger U$;
P(12) $|(T^*)^\dagger| = |T|^\dagger$;
P(13) $T \geq 0 \Leftrightarrow T^\dagger \geq 0$;
P(14) $U^*|T^*|$ and $U^*|T^*|^\dagger$ are the polar decompositions of T^* and T^\dagger .

Let f be a bounded Borel real-valued function defined in an interval $\mathcal{I} \subseteq \mathbb{R}$. If $T \in B(H)$ is a self-adjoint operator, then by $f(T)$ we mean the self-adjoint operator $\int_{-\infty}^{+\infty} f(\lambda)dE_\lambda$ where E_λ is the spectral resolution of identity corresponding to T . The restriction of f to the set of all self-adjoint operators is called an operator function. For example, for each $q > 0$, $f(x) = x^q$ is an operator function. In addition, in this case, if U is any unitary operator, then $f(U^*TU) = U^*f(T)U$. For more details see [3, 4].

Lemma 2.1. *Let $T \in CR(H)$. Then the following assertions hold.*

- (i) $(|T|^\dagger)^q = U^*(|T^*|^\dagger)^qU$, for each $q > 0$.
(ii) T^\dagger is quasinormal if and only if $U^*|T^*|^\dagger = |T^*|^\dagger U^*$.

Proof. (i) By P(3), P(10), P(11) and P(14) we have

$$\begin{aligned} (|T|^\dagger)^2 &= |(T^*)^\dagger|^2 = |(T^\dagger)^*|^2 = T^\dagger(T^\dagger)^* \\ &= U^*|T^*|^\dagger|T^*|^\dagger U \\ &= U^*|T^*|^\dagger U U^*|T^*|^\dagger U \\ &= (U^*|T^*|^\dagger U)^2. \end{aligned}$$

Since for each $q > 0$, $f(x) = x^{\frac{q}{2}}$ is an operator function, we obtain $(|T|^\dagger)^q = U^*(|T^*|^\dagger)^qU$.

(ii) It is a classical fact that T is quasinormal if and only if $U|T| = |T|U$ (see for example [12, Theorem 3]). Now, the desired conclusion follows from this and P(14). \square

Theorem 2.2. *Let $T \in B(H)$ be onto. Then the following statements are equivalent:*

- (i) T^* is quasinormal.

(ii) T^\dagger is quasinormal.

Moreover, if one of the above statements hold then

$$(iii) [(T^\dagger)^* T^\dagger, (T^\dagger)^* + T^\dagger] = 0.$$

Proof. (i) \Leftrightarrow (ii) Since $|T^*| |T^*|^\dagger |T^*| = |T^*|$, then we have

$$\begin{aligned} T^* \text{ is quasinormal} &\Leftrightarrow U^* |T^*| = |T^*| U^* \\ &\Leftrightarrow U^* |T^*| |T^*|^\dagger |T^*| = |T^*| |T^*|^\dagger |T^*| U^* \\ &\Leftrightarrow |T^*| U^* |T^*|^\dagger |T^*| = |T^*| |T^*|^\dagger U^* |T^*| \\ &\Leftrightarrow |T^*| (U^* |T^*|^\dagger - |T^*|^\dagger U^*) |T^*| = 0. \end{aligned}$$

By hypothesis, $\mathcal{N}(|T^*|) = \mathcal{N}(T^*) = \{0\}$. Hence $(U^* |T^*|^\dagger - |T^*|^\dagger U^*) |T^*| = 0$, and so $U^* |T^*|^\dagger = |T^*|^\dagger U^*$ on $\overline{R(|T^*|^\dagger)}$. On the other hand, $U^* |T^*|^\dagger = |T^*|^\dagger U^*$ on $N(|T^*|^\dagger) = N(U^*)$. Thus, $U^* |T^*|^\dagger = |T^*|^\dagger U^*$ on H . Consequently, by Lemma 2.1(ii), (i) \Leftrightarrow (ii) holds.

Now, it is easy to check that

$$\begin{aligned} [(T^\dagger)^* T^\dagger, (T^\dagger)^* + T^\dagger] &= [(|T^*|^\dagger)^2, |T^*|^\dagger U + U^* |T^*|^\dagger] \\ &= \{(|T^*|^\dagger)^2 U^* |T^*|^\dagger - U^* (|T^*|^\dagger)^3\} + |T^*|^\dagger \{(|T^*|^\dagger)^2 U - U (|T^*|^\dagger)^2\}. \end{aligned}$$

If (ii) is holds, then by Lemma 2.1(ii) we obtain

$$\begin{aligned} (|T^*|^\dagger)^2 U^* |T^*|^\dagger &= U^* (|T^*|^\dagger)^3; \\ (|T^*|^\dagger)^2 U &= U (|T^*|^\dagger)^2. \end{aligned}$$

Thus, $[(T^\dagger)^* T^\dagger, (T^\dagger)^* + T^\dagger] = 0$. \square

For more details and applications on condition (iii) in Theorem 2.2 see [16].

Lemma 2.3. *If $T \in \text{CR}(H)$ is binormal, then $(\widetilde{T^*})^\dagger = (|T^*|^\dagger)^{\frac{1}{2}} U (|T^*|^\dagger)^{\frac{1}{2}}$.*

Proof. Since T is binormal, then we obtain from direct computations that

$$\begin{aligned} \widetilde{T^*} (\widetilde{T^*})^\dagger \widetilde{T^*} &= |T^*|^{\frac{1}{2}} U^* |T^*|^{\frac{1}{2}} (|T^*|^\dagger)^{\frac{1}{2}} U (|T^*|^\dagger)^{\frac{1}{2}} |T^*|^{\frac{1}{2}} U^* |T^*|^{\frac{1}{2}} \\ &= |T^*|^{\frac{1}{2}} U^* P_{R(|T^*|)} U P_{R(|T^*|)} U^* |T^*|^{\frac{1}{2}} \quad \text{by P(4)} \\ &= |T^*|^{\frac{1}{2}} U^* U P_{R(|T^*|)} U^* |T^*|^{\frac{1}{2}} \quad \text{by P(6)} \\ &= |T^*|^{\frac{1}{2}} P_{R(|T|)} P_{R(|T^*|)} U^* |T^*|^{\frac{1}{2}} \quad \text{by P(7)} \\ &= |T^*|^{\frac{1}{2}} P_{R(|T^*|)} P_{R(|T|)} U^* |T^*|^{\frac{1}{2}} \quad \text{by P(5)} \\ &= |T^*|^{\frac{1}{2}} U^* |T^*|^{\frac{1}{2}} = \widetilde{T^*} \quad \text{by P(4), P(6)}. \end{aligned}$$

Also,

$$\begin{aligned}
(\widetilde{T^*})^\dagger \widetilde{T^*} (\widetilde{T^*})^\dagger &= (|T^*|^\dagger)^{\frac{1}{2}} U (|T^*|^\dagger)^{\frac{1}{2}} |T^*|^{\frac{1}{2}} U^* |T^*|^{\frac{1}{2}} (|T^*|^\dagger)^{\frac{1}{2}} U (|T^*|^\dagger)^{\frac{1}{2}} \\
&= (|T^*|^\dagger)^{\frac{1}{2}} U P_{R(|T^*|)} U^* P_{R(|T^*|)} U (|T^*|^\dagger)^{\frac{1}{2}} && \text{by P(4)} \\
&= (|T^*|^\dagger)^{\frac{1}{2}} U P_{R(|T^*|)} U^* U (|T^*|^\dagger)^{\frac{1}{2}} && \text{by P(6)} \\
&= (|T^*|^\dagger)^{\frac{1}{2}} U P_{R(|T^*|)} P_{R(|T|)} (|T^*|^\dagger)^{\frac{1}{2}} && \text{by P(7)} \\
&= (|T^*|^\dagger)^{\frac{1}{2}} U P_{R(|T|)} P_{R(|T^*|)} (|T^*|^\dagger)^{\frac{1}{2}} && \text{by P(5)} \\
&= (|T^*|^\dagger)^{\frac{1}{2}} P_{R(|T^*|)} U P_{R(|T^*|)} (|T^*|^\dagger)^{\frac{1}{2}} && \text{by P(6)} \\
&= (|T^*|^\dagger)^{\frac{1}{2}} U (|T^*|^\dagger)^{\frac{1}{2}} = (\widetilde{T^*})^\dagger && \text{by P(4)}.
\end{aligned}$$

Similar computations show that

$$(\widetilde{T^*})^\dagger \widetilde{T^*} = U P_{R(|T^*|)} P_{R(|T|)} U^*$$

and

$$\widetilde{T^*} (\widetilde{T^*})^\dagger = P_{R(|T|)} P_{R(|T^*|)}.$$

Hence, $(\widetilde{T^*})^\dagger \widetilde{T^*}$ and $\widetilde{T^*} (\widetilde{T^*})^\dagger$ are self-adjoint operators. This completes the proof. \square

Note that if $T \in \text{CR}(\mathbb{H})$ is binormal, then Lemma 2.3 shows that $\widetilde{T^*}$ and so \widetilde{T} have closed range. Moreover, in this case, we have $\widetilde{T}^\dagger = (|T|^\dagger)^{\frac{1}{2}} U^* (|T|^\dagger)^{\frac{1}{2}}$.

Theorem 2.4. *Let $T \in B(H)$ be onto and binormal. Then T^* is quasinormal if and only if $(T^\dagger)^* = (\widetilde{T^*})^\dagger$.*

Proof. By [21, Theorem 10] and Theorem 2.2, T^* is quasinormal if and only if $T^\dagger = \widetilde{T}^\dagger$. Now, taking adjoint of both sides and using Lemma 2.3 and P(1), we obtain $(T^\dagger)^* = (\widetilde{T^*})^\dagger$. \square

In the following, we concentrate on the polar decomposition of $(\widetilde{T})^\dagger$. We require the following lemma.

Lemma 2.5. *(i) [11, Corollary 1] Let $T = U|T|$ and $S = V|S|$ be the polar decompositions. If T and S are doubly commutative (i.e., $[T, S] = [T, S^*] = 0$), then $TS = UV|TS|$.*

(ii) [22, Proposition 3.9] Let $T = U|T|$ be the polar decomposition of a binormal operator T . Then $\widetilde{T} = U^ U U |\widetilde{T}|$ is also the polar decomposition of \widetilde{T} .*

(iii) [22, Theorem 2.1] Let $T = U|T|$ and $|T|^{\frac{1}{2}} |T^|^{\frac{1}{2}} = V ||T|^{\frac{1}{2}} |T^*|^{\frac{1}{2}}$ be the polar decompositions. Then $\widetilde{T} = V U |\widetilde{T}|$ is also the polar decomposition.*

Theorem 2.6. *Let $T = U|T| \in \text{CR}(\mathbb{H})$ and $(|T|^\dagger)^{\frac{1}{2}}(|T^*|^\dagger)^{\frac{1}{2}} = V(|T|^\dagger)^{\frac{1}{2}}(|T^*|^\dagger)^{\frac{1}{2}}$ be the polar decompositions. If T is binormal, then $(\tilde{T})^\dagger = U^*V|(\tilde{T})^\dagger|$ is also the polar decomposition.*

Proof. (i) First we show that $(\tilde{T})^\dagger = U^*V|(\tilde{T})^\dagger|$.

$$\begin{aligned}
U^*V|(\tilde{T})^\dagger| &= U^*V((\tilde{T})^\dagger)^*(\tilde{T})^\dagger \\
&= U^*V(|T|^\dagger)^{\frac{1}{2}}U(|T|^\dagger)U^*(|T|^\dagger)^{\frac{1}{2}} && \text{by Lemma 2.3} \\
&= U^*V(|T|^\dagger)^{\frac{1}{2}}UU^*(|T^*|^\dagger)UU^*(|T|^\dagger)^{\frac{1}{2}} && \text{by Lemma 2.1(i)} \\
&= U^*V(|T|^\dagger)^{\frac{1}{2}}|T^*|^\dagger(|T|^\dagger)^{\frac{1}{2}} && \text{by P(8),P(10)} \\
&= U^*V(|T|^\dagger)^{\frac{1}{2}}(|T^*|^\dagger)^{\frac{1}{2}} \\
&= U^*(|T|^\dagger)^{\frac{1}{2}}(|T^*|^\dagger)^{\frac{1}{2}} \\
&= U^*(|T^*|^\dagger)^{\frac{1}{2}}(|T|^\dagger)^{\frac{1}{2}} \\
&= U^*(|T^*|^\dagger)^{\frac{1}{2}}UU^*(|T|^\dagger)^{\frac{1}{2}} \\
&= (|T|^\dagger)^{\frac{1}{2}}U^*(|T|^\dagger)^{\frac{1}{2}} = (\tilde{T})^\dagger.
\end{aligned}$$

Now, we claim that $N((\tilde{T})^\dagger) = N(U^*V)$. Since T is binormal, then it is easy to check that T^\dagger is binormal. Thus $N((|T^*|^\dagger)^{\frac{1}{2}}(|T|^\dagger)^{\frac{1}{2}}) = N((|T|^\dagger)^{\frac{1}{2}}(|T^*|^\dagger)^{\frac{1}{2}}) = N(V)$. Then we have

$$\begin{aligned}
U^*Vx = 0 &\Leftrightarrow U^*(|T|^\dagger)^{\frac{1}{2}}(|T^*|^\dagger)^{\frac{1}{2}}x = 0 \\
&\Leftrightarrow U^*(|T^*|^\dagger)^{\frac{1}{2}}(|T|^\dagger)^{\frac{1}{2}}x = 0 \\
&\Leftrightarrow U^*(|T^*|^\dagger)^{\frac{1}{2}}UU^*(|T|^\dagger)^{\frac{1}{2}}x = 0 && \text{by P(8)} \\
&\Leftrightarrow (|T|^\dagger)^{\frac{1}{2}}U^*(|T|^\dagger)^{\frac{1}{2}}x = 0 && \text{by Lemma 2.1(i)} \\
&\Leftrightarrow (\tilde{T})^\dagger x = 0.
\end{aligned}$$

Lastly, we prove that U^*V is partial isometry. Since $(|T|^\dagger)^{\frac{1}{2}} = U^*U(|T|^\dagger)^{\frac{1}{2}}$ and $(|T^*|^\dagger)^{\frac{1}{2}} = UU^*(|T^*|^\dagger)^{\frac{1}{2}}$ are the polar decompositions of $(|T|^\dagger)^{\frac{1}{2}}$ and $(|T^*|^\dagger)^{\frac{1}{2}}$, respectively then by Lemma 2.5(i) we have

$$(|T|^\dagger)^{\frac{1}{2}}(|T^*|^\dagger)^{\frac{1}{2}} = (|T^*|^\dagger)^{\frac{1}{2}}(|T|^\dagger)^{\frac{1}{2}} = UU^*U^*U(|T^*|^\dagger)^{\frac{1}{2}}(|T|^\dagger)^{\frac{1}{2}}.$$

Then by the uniqueness of the polar decomposition we get that $V = UU^*U^*U$. It

follows that

$$\begin{aligned}
(U^*V)(U^*V)^*(U^*V) &= (U^*UU^*U^*U)(U^*UU^*U^*U)^*(U^*UU^*U^*U) \\
&= P_{R(|T|)}U^*P_{R(|T|)}UP_{R(|T|)}U^*P_{R(|T|)} && \text{by P(7)} \\
&= U^*P_{R(|T|)}UU^*P_{R(|T|)} && \text{by P(6)} \\
&= U^*P_{R(|T|)}P_{R(|T^*|)}P_{R(|T|)} && \text{by P(6)} \\
&= U^*P_{R(|T^*|)}P_{R(|T|)} && \text{by P(5)} \\
&= P_{R(|T|)}U^*P_{R(|T|)} && \text{by P(6)} \\
&= U^*UU^*U^*U = U^*V.
\end{aligned}$$

This completes the proof. \square

Corollary 2.7. *Let $T \in \text{CR}(\mathbb{H})$ be binormal and let $T^\dagger = U^*|T^*|^\dagger$ and $(|T|^\dagger)^{\frac{1}{2}}(|T^*|^\dagger)^{\frac{1}{2}} = V(|T|^\dagger)^{\frac{1}{2}}(|T^*|^\dagger)^{\frac{1}{2}}$ be the polar decompositions. Then the following statements are hold:*

- (i) $\widetilde{T}^\dagger = UU^*U^*|\widetilde{T}^\dagger|$ is the polar decomposition.
- (ii) $(\widetilde{T})^\dagger = U^*U^*U|(\widetilde{T})^\dagger|$ is the polar decomposition.

Proof. (i) Since T is binormal, T^\dagger is binormal. Now, the desired conclusion follows by Lemma 2.5(iii).

(ii) Recall that $(|T|^\dagger)^{\frac{1}{2}} = U^*U(|T|^\dagger)^{\frac{1}{2}}$ and $(|T^*|^\dagger)^{\frac{1}{2}} = UU^*(|T^*|^\dagger)^{\frac{1}{2}}$ are the polar decompositions of $(|T|^\dagger)^{\frac{1}{2}}$ and $(|T^*|^\dagger)^{\frac{1}{2}}$, respectively. Then by Lemma 2.5(i) we obtain

$$(|T|^\dagger)^{\frac{1}{2}}(|T^*|^\dagger)^{\frac{1}{2}} = (|T^*|^\dagger)^{\frac{1}{2}}(|T|^\dagger)^{\frac{1}{2}} = UU^*U^*U(|T^*|^\dagger)^{\frac{1}{2}}(|T|^\dagger)^{\frac{1}{2}}.$$

Thus, by Theorem 2.6, $(\widetilde{T})^\dagger = U^*UU^*U^*U|(\widetilde{T})^\dagger| = U^*U^*U|(\widetilde{T})^\dagger|$. \square

In [19], Yamazaki introduce the notion of the $*$ -Aluthge transformation $\widetilde{T}^{(*)}$ of T by setting $\widetilde{T}^{(*)} = |T^*|^{\frac{1}{2}}U|T^*|^{\frac{1}{2}}$. Like this notion we introduce \dagger -Aluthge transformation $\widetilde{T}^{(\dagger)}$ of T by setting $\widetilde{T}^{(\dagger)} = (\widetilde{T}^\dagger)^\dagger$. Similar computations show that $\widetilde{T}^{(\dagger)} = \widetilde{T}^{(*)}$, whenever $T \in \text{CR}(\mathbb{H})$ is binormal.

Proposition 2.8. *Let $T \in \text{CR}(\mathbb{H})$. Then the following statements hold.*

- (i) *If T is self-adjoint, then $W(T) \subseteq W(T^\dagger)W(T^2)$.*
- (ii) *If T is onto and T^* is quasinormal, then $W(\widetilde{T}^\dagger) = W(T^\dagger) \subseteq W(U^*)W(|T^*|^\dagger)$.*
- (iii) *If T is binormal, then $W((\widetilde{T})^{(\dagger)}) \subseteq W(U)W(|T^*|)$.*

Proof. (i) Let $x \in H$ with $\|x\| = 1$. Then we get that

$$\begin{aligned}\langle Tx, x \rangle &= \langle TT^\dagger Tx, x \rangle = \langle T^\dagger Tx, Tx \rangle \\ &= \left\langle T^\dagger \frac{Tx}{\|Tx\|}, \frac{Tx}{\|Tx\|} \right\rangle \langle Tx, Tx \rangle.\end{aligned}$$

It follows that $\langle Tx, x \rangle \in W(T^\dagger)W(T^2)$, for each $x \in H$ with $\|x\| = 1$.

(ii) By Theorem 2.2 and Lemma 2.1, T^\dagger is quasinormal and so $U^*|T^*|^\dagger = |T^*|^\dagger U^*$. It follows that $U^*(|T^*|^\dagger)^{\frac{1}{2}} = (|T^*|^\dagger)^{\frac{1}{2}}U^*$. Then by P(1) and P(14) we have

$$\begin{aligned}\langle T^\dagger x, x \rangle &= \langle U^*|T^*|^\dagger x, x \rangle = \langle \widetilde{T}^\dagger x, x \rangle \\ &= \langle U^*(|T^*|^\dagger)^{\frac{1}{2}}x, (|T^*|^\dagger)^{\frac{1}{2}}x \rangle \\ &= \left\langle U^* \frac{(|T^*|^\dagger)^{\frac{1}{2}}x}{\|(|T^*|^\dagger)^{\frac{1}{2}}x \|}, \frac{(|T^*|^\dagger)^{\frac{1}{2}}x}{\|(|T^*|^\dagger)^{\frac{1}{2}}x \|} \right\rangle \langle (|T^*|^\dagger)x, x \rangle,\end{aligned}$$

for each $x \in H$ with $\|x\| = 1$.

(iii) Direct replacement shows that

$$\begin{aligned}\langle \widetilde{T}^{(\dagger)}x, x \rangle &= \langle |T^*|^{\frac{1}{2}}U|T^*|^{\frac{1}{2}}x, x \rangle \\ &= \langle U|T^*|^{\frac{1}{2}}x, |T^*|^{\frac{1}{2}}x \rangle \\ &= \left\langle U \frac{|T^*|^{\frac{1}{2}}x}{\||T^*|^{\frac{1}{2}}x \|}, \frac{|T^*|^{\frac{1}{2}}x}{\||T^*|^{\frac{1}{2}}x \|} \right\rangle \langle |T^*|^{\frac{1}{2}}x, x \rangle,\end{aligned}$$

for each $x \in H$ with $\|x\| = 1$. This completes the proof. \square

Proposition 2.9. *Let $T \in \text{CR}(H)$. Then the following assertions hold.*

(i) *Let $T \in \text{CR}(H)$ be self-adjoint. Then $\omega(T) \leq \omega(T^\dagger)\|T\|^2$.*

(ii) *If T is onto and T^* be quasinormal, then*

$$\omega(\widetilde{T}^\dagger) = \omega(T^\dagger) \leq \omega(U^*)\||T^*|^\dagger\|.$$

Proof. (i) Since T is self-adjoint and $T = TT^\dagger T$, we have

$$\begin{aligned}\omega(T) &= \sup_{\|x\|=1} |\langle TT^\dagger Tx, x \rangle| = \sup_{\|x\|=1} |\langle T^\dagger Tx, Tx \rangle| \\ &= \sup_{\|x\|=1} \left| \left\langle T^\dagger \frac{Tx}{\|Tx\|}, \frac{Tx}{\|Tx\|} \right\rangle \right| \|Tx\|^2 \\ &\leq \omega(T^\dagger)\|T\|^2.\end{aligned}$$

(ii) Since T^* is quasinormal then by Theorem 2.2, we obtain

$$\begin{aligned}\omega(T^\dagger) &= \sup_{\|x\|=1} |\langle T^\dagger x, x \rangle| \\ &= \sup_{\|x\|=1} |\langle U^* |T^*|^\dagger x, x \rangle| \\ &= \sup_{\|x\|=1} |\langle (|T^*|^\dagger)^{\frac{1}{2}} U^* (|T^*|^\dagger)^{\frac{1}{2}} x, x \rangle| = \omega(\widetilde{T}^\dagger).\end{aligned}$$

Thus,

$$\begin{aligned}\omega(T^\dagger) &= \sup_{\|x\|=1} |\langle T^\dagger x, x \rangle| \\ &= \sup_{\|x\|=1} |\langle U^* |T^*|^\dagger x, x \rangle| \\ &= \sup_{\|x\|=1} |\langle U^* (|T^*|^\dagger)^{\frac{1}{2}} x, (|T^*|^\dagger)^{\frac{1}{2}} x \rangle| \\ &= \sup_{\|x\|=1} |\langle U^* \frac{(|T^*|^\dagger)^{\frac{1}{2}} x}{\|(|T^*|^\dagger)^{\frac{1}{2}} x\|}, \frac{(|T^*|^\dagger)^{\frac{1}{2}} x}{\|(|T^*|^\dagger)^{\frac{1}{2}} x\|} \rangle| \| |T^*|^\dagger x \| \\ &\leq \omega(U^*) \| |T^*|^\dagger \|.\end{aligned}$$

□

Proposition 2.10. *Let $T \in \text{CR}(\mathbb{H})$. Then the following assertions hold.*

$$(i) \omega(\widetilde{T}^\dagger) \leq \omega(|T^*|) \leq \| |T^*| \|.$$

$$(ii) \text{ If } T \text{ is binormal, then } \omega(\widetilde{T}^\dagger) = \omega(T^\dagger) \text{ and } \|\widetilde{T}^\dagger\| = \|T^\dagger\|.$$

Proof. The first part is easily follow from Proposition 2.8. For the second part, since

$$\begin{aligned}\widetilde{T}^\dagger &= (|T|^\dagger)^{\frac{1}{2}} U^* (|T|^\dagger)^{\frac{1}{2}} && \text{by Lemma 2.3} \\ &= U^* (|T^*|^\dagger)^{\frac{1}{2}} U U^* U^* (|T^*|^\dagger)^{\frac{1}{2}} U && \text{by Lemma 2.1(i)} \\ &= U^* (|T^*|^\dagger)^{\frac{1}{2}} U^* (|T^*|^\dagger)^{\frac{1}{2}} U && \text{by P(8)} \\ &= U^* |T^\dagger|^{\frac{1}{2}} U^* |T^\dagger|^{\frac{1}{2}} U && \text{by P(3)} \\ &= U^* \widetilde{T}^\dagger U && \text{by P(1),}\end{aligned}$$

then

$$\begin{aligned}\omega(\widetilde{T}^\dagger) &= \sup_{\|x\|=1} |\langle U^* \widetilde{T}^\dagger U x, x \rangle| = \sup_{\|x\|=1} |\langle \widetilde{T}^\dagger U x, U x \rangle| \\ &= \sup_{\|x\|=1} |\langle \widetilde{T}^\dagger \frac{U x}{\|U x\|}, \frac{U x}{\|U x\|} \rangle| \|U x\|^2 \leq \omega(\widetilde{T}^\dagger).\end{aligned}$$

On the other hand, since

$$\begin{aligned}
\widetilde{T}^\dagger &= (|T^*|^\dagger)^{\frac{1}{2}} U^* (|T^*|^\dagger)^{\frac{1}{2}} && \text{by P(1)} \\
&= (|T^*|^\dagger)^{\frac{1}{2}} U U^* U^* (|T^*|^\dagger)^{\frac{1}{2}} && \text{by P(8)} \\
&= U (|T|^\dagger)^{\frac{1}{2}} U^* (|T|^\dagger)^{\frac{1}{2}} U^* && \text{by P(9)} \\
&= U \widetilde{T}^\dagger U^* && \text{by Lemma 2.3,}
\end{aligned}$$

then

$$\begin{aligned}
\omega(\widetilde{T}^\dagger) &= \sup_{\|x\|=1} |\langle U \widetilde{T}^\dagger U^* x, x \rangle| = \sup_{\|x\|=1} |\langle \widetilde{T}^\dagger U^* x, U^* x \rangle| \\
&= \sup_{\|x\|=1} |\langle \widetilde{T}^\dagger \frac{U^* x}{\|U^* x\|}, \frac{U^* x}{\|U^* x\|} \rangle| \|U^* x\|^2 \leq \omega(\widetilde{T}^\dagger).
\end{aligned}$$

Moreover, since $\widetilde{T}^\dagger = U^* \widetilde{T}^\dagger U$ and $\widetilde{T}^\dagger = U^* \widetilde{T}^\dagger U$, we obtain $\|\widetilde{T}^\dagger\| = \|\widetilde{T}^\dagger\|$. \square

Lemma 2.11. [7, Theorem 2.8] *If A be an arbitrary operator and B is normal. Then $Sp(AB) = Sp(BA)$.*

Proposition 2.12. *Let $T \in CR(H)$. Then the following statements hold.*

(i) $(T^*)^\dagger = U|T|^\dagger$ is the polar decomposition.

(ii) If T is binormal, then $Sp(T^\dagger) = Sp(\widetilde{T}^\dagger) = Sp(\widetilde{T}^\dagger)$.

(iii) If T is binormal and $\lambda > 0$, then $\lambda \in Sp(T) \Leftrightarrow \lambda \in Sp((\widetilde{T}^\dagger)^\dagger)$.

Proof. (i) Since $U^*|T^*|^\dagger$ is the unique polar decomposition of T^\dagger , then $N(U^*) = N(|T^*|^\dagger)$, and so $N(U^*U) = N(|T^*|^\dagger U)$. Now, by Lemma 2.1(i), P(11) and P(12) we have

$$(2.1) \quad (T^*)^\dagger = (T^\dagger)^* = |T^*|^\dagger U = U U^* |T^*|^\dagger U = U |T|^\dagger;$$

$$N(U) = N(U^*U) = N(|T^*|^\dagger U) = N((T^*)^\dagger) = N(|(T^\dagger)^*|) = N(|T|^\dagger).$$

Therefore, $(T^*)^\dagger = U|T|^\dagger$ is the unique polar decomposition.

(ii) By P(14), $(|T|^\dagger)^{\frac{1}{2}} \geq 0$ and hence it is normal. Thus, by (2.1) and Lemma 2.11 we have

$$\begin{aligned}
Sp(T^\dagger) &= Sp(|T|^\dagger U^*) = Sp((|T|^\dagger)^{\frac{1}{2}} (|T|^\dagger)^{\frac{1}{2}} U^*) \\
&= Sp((|T|^\dagger)^{\frac{1}{2}} U^* (|T|^\dagger)^{\frac{1}{2}}) = Sp(\widetilde{T}^\dagger).
\end{aligned}$$

Similarly, we get that

$$\begin{aligned}
Sp(\widetilde{T}^\dagger) &= Sp((|T^*|^\dagger)^{\frac{1}{2}} U^* (|T^*|^\dagger)^{\frac{1}{2}}) \\
&= Sp(U^* (|T^*|^\dagger)^{\frac{1}{2}} (|T^*|^\dagger)^{\frac{1}{2}}) = Sp(T^\dagger).
\end{aligned}$$

(iii) If $\lambda > 0$, then by Lemma 2.3 and part (ii) we have

$$\begin{aligned}\lambda \in Sp(T) &\Leftrightarrow \lambda^{-1} \in Sp(T^\dagger) \Leftrightarrow \lambda^{-1} \in Sp(\widetilde{T}^\dagger) \\ &\Leftrightarrow \lambda \in Sp((\widetilde{T}^\dagger)^\dagger) = Sp((\widetilde{T})^{(\dagger)}).\end{aligned}$$

□

Recall that an operator $T \in CR(H)$ is an *EP* operator if and only if $TT^\dagger = T^\dagger T$ [8]. If $T = T^\dagger$, then $T^3 = TT^\dagger T = T$ and hence $T^{2n+1} = T$ for all $n \in \mathbb{N}$. On the other hand, if $TT^\dagger = T^\dagger T$ and $T^{2n+1} = T$, then for $n = 3$ we obtain $T^\dagger = T^\dagger T T^\dagger = T^\dagger T^3 T^\dagger$ and thus

$$T^\dagger = T^\dagger T T T T^\dagger = T T^\dagger T T T^\dagger = T T^\dagger T T^\dagger T = T T^\dagger T = T.$$

Now, let $T^{2n+3} = T$. Since T is an *EP* operator, then

$$T^\dagger = T^\dagger T T^\dagger = T^\dagger T^{2n+3} T^\dagger = T^\dagger T^{2n+1} T^\dagger T^2 = T^\dagger T^2 = T T^\dagger T = T.$$

These observations establish the following proposition.

Proposition 2.13. *Let $T \in CR(H)$ and $n \in \mathbb{N}$. Then the following statements hold.*

(i) *If $T = T^\dagger$, then $T = T^{2n+1}$.*

(ii) *If $T = T^{2n+1}$ and T is an *EP* operator, then $T = T^\dagger$.*

3. APPLICATIONS TO THE LAMBERT MULTIPLICATION OPERATORS

Let (X, Σ, μ) be a complete σ -finite measure space. For any complete σ -finite subalgebra $\mathcal{A} \subseteq \Sigma$ the Hilbert space $L^2(X, \mathcal{A}, \mu|_{\mathcal{A}})$ is abbreviated to $L^2(\mathcal{A})$ where $\mu|_{\mathcal{A}}$ is the restriction of μ to \mathcal{A} . We denote the linear space of all complex-valued Σ -measurable functions on X by $L^0(\Sigma)$. The support of a measurable function f is defined by $\sigma(f) = \{x \in X : f(x) \neq 0\}$. All sets and functions statements are to be interpreted as being valid almost everywhere with respect to μ . For each non-negative $f \in L^0(\Sigma)$ or $f \in L^2(\Sigma)$, by the Radon-Nikodym theorem, there exists a unique \mathcal{A} -measurable function $E^{\mathcal{A}}(f)$ such that

$$\int_A f d\mu = \int_A E^{\mathcal{A}}(f) d\mu,$$

where A is any \mathcal{A} -measurable set for which $\int_A f d\mu$ exists. Now associated with every complete σ -finite subalgebra $\mathcal{A} \subseteq \Sigma$, the mapping $E^{\mathcal{A}} : L^2(\Sigma) \rightarrow L^2(\mathcal{A})$

uniquely defined by the assignment $f \mapsto E^{\mathcal{A}}(f)$, is called the conditional expectation operator with respect to \mathcal{A} . Put $E = E^{\mathcal{A}}$. The mapping E is a linear orthogonal projection onto $L^2(\mathcal{A})$. Note that $\mathcal{D}(E)$, the domain of E , contains $L^2(\Sigma) \cup \{f \in L^0(\Sigma) : f \geq 0\}$. For more details on the properties of E see [14, 18].

We shall always take $u \in L^0(\Sigma)$ for which $uf \in \mathcal{D}(E)$ for all $f \in L^2(\Sigma)$. In other words, EM_u is a well-defined operator on $L^2(\Sigma)$. The mapping $T : L^2(\Sigma) \rightarrow L^2(\Sigma)$ defined by $T(f) = wE(uf)$ is called Lambert multiplication operator. For other important properties of T see ([5, 6, 15]). By [15, Proposition 2.1(b)], EM_u is bounded on $L^2(\Sigma)$ if and only if $E(|u|^2) \in L^\infty(\mathcal{A})$. In this case $\|EM_u\| = \|E(|u|^2)\|_\infty^{1/2}$. Now, let $f \in L^2(\Sigma)$. Then

$$\begin{aligned} \|Tf\|^2 &= \int E(|w|^2)|E(uf)|^2 d\mu = \int |E(u(E(|w|^2))^{\frac{1}{2}}f)|^2 d\mu \\ (3.2) \qquad &= \int |E(M_v f)|^2 d\mu = \|EM_v f\|^2, \end{aligned}$$

where $v := u(E(|w|^2))^{\frac{1}{2}}$. It follows that $T = M_w EM_u$ is bounded on $L^2(\Sigma)$ if and only if $E(|w|^2)E(|u|^2) \in L^\infty(\mathcal{A})$, and in this case $\|T\| = \|E(|w|^2)^{1/2}E(|u|^2)^{1/2}\|_\infty$. Now, let $0 \leq u \in L^0(\Sigma)$ and let $E(u) \geq \delta$ on $S := \sigma(E(u))$. Note that $L^2(\Sigma) = L^2(S) \oplus L^2(S^c)$, where $S^c = X \setminus S$, $L^2(S) = L^2(S, \Sigma_S, \mu|_S)$ and $\Sigma_S = \{A \cap S : A \in \Sigma\}$. We claim that $T_1 := EM_u$ has closed range. To this end let $f_n, g \in L^2(\Sigma)$ with $\|g\|_2 > 0$ and $T_1 f_n \rightarrow g$ in $L^2(\Sigma)$. Since $L^2(S^c) \subseteq N(T)$, then $g = 0$ on S^c and hence $T_1 f_n \rightarrow \chi_S g$ in $L^2(\Sigma)$. But $\chi_S g = EM_u(\frac{\chi_S g}{E(u)})$, because $g \in L^2(\mathcal{A})$ and

$$\|\frac{\chi_S g}{E(u)}\|_2 \leq \frac{1}{\delta} \|g\|_2.$$

It follows that $g = \chi_{S^c} g + \chi_S g = 0 + E(\frac{\chi_S u g}{E(u)}) \in R(T_1)$, and so T_1 has closed range. By (3.1), $T \in B(L^2(\Sigma))$ has closed range if and only if $T_1 \in B(L^2(\Sigma))$ has closed range. These observations establish the following proposition.

Proposition 3.14. *Let $T : L^2(\Sigma) \rightarrow L^0(\Sigma)$ defined by $T = M_w EM_u$ is a Lambert multiplication operator.*

(i) $T \in B(L^2(\Sigma))$ if and only if $E(|w|^2)E(|u|^2) \in L^\infty(\mathcal{A})$, and in this case $\|T\| = \|E(|w|^2)E(|u|^2)\|_\infty^{1/2}$.

(ii) Let $T \in B(L^2(\Sigma))$, $0 \leq u \in L^0(\Sigma)$ and $v = u(E(|w|^2))^{\frac{1}{2}}$. If $E(v) \geq \delta$ on $\sigma(v)$, then T has closed range.

In what follows, since for each $u \geq 0$, $\sigma(u) \subseteq \sigma(E(u^2))$, we use the notational convention of $\frac{u}{E(u)}$ for $\frac{u}{E(u)} \chi_{\sigma(u)}$. From now on, we assume that $u, w \in \mathcal{D}(E)$ are

non-negative, $S = \sigma(E(u^2)) = \sigma(E(u))$ and $T = M_w E M_u \in \text{CR}(L^2(\Sigma))$.

Let B, C be bounded positive operators on H such that $BC = CB$. Put $A = BC$. Since $f(x) = x^p$ is an operator function, we obtain $A^p = B^p C^p$ for each $p > 0$. In particular, take $B = M_\nu$ and $C = M_{\bar{\omega}} E M_\omega$, where $0 \leq \nu \in L^0(\mathcal{A})$ and $\omega \in L^0(\Sigma)$. A direct computations shows that $C^p = M_{\omega E(|\omega|^2)^{p-1}} E M_\omega$. Consequently, we have the following lemma.

Lemma 3.15. *Let $0 \leq \nu \in L^0(\mathcal{A})$, $\omega \in L^0(\Sigma)$ and let $A := M_{\nu \bar{\omega}} E M_\omega \in B(L^2(\Sigma))$. Then for each $p \in (0, \infty)$, $A^p = M_{\nu^p \bar{\omega} E(|\omega|^2)^{p-1}} E M_\omega$.*

Put

$$(3.3) \quad A(f) = \frac{u \chi_G}{E(u^2) E(w^2)} E(wf), \quad f \in L^2(\Sigma), \quad G = \sigma(E(w)).$$

Then by Proposition 3.1, $A \in B(L^2(\Sigma))$. Also, it is easy to check that

$$TAT = T, \quad ATA = A, \quad (TA)^* = TA, \quad (AT)^* = AT.$$

Thus, $A = T^\dagger = M_{\frac{\chi_{S \cap G}}{E(u^2) E(w^2)}} T^*$ and hence A has closed range.

Now, we concentrate on the parts of the polar decomposition T , T^\dagger and their Aluthge transformations. Let $f \in L^2(\Sigma)$. Then we can obtain from direct computations that

$$\begin{aligned} |T|^2(f) &= \chi_S u E(w^2) E(uf); \\ |T|(f) &= u (E(u^2))^{-\frac{1}{2}} (E(w^2))^{\frac{1}{2}} E(uf) && \text{by Lemma 3.2;} \\ |T|^{\frac{1}{2}}(f) &= u (E(u^2))^{-\frac{3}{4}} (E(w^2))^{\frac{1}{4}} E(uf) && \text{by Lemma 3.2;} \\ U(f) &= \chi_S w (E(u^2))^{-\frac{1}{2}} (E(w^2))^{-\frac{1}{2}} E(uf) && \text{because } U|T| = T. \end{aligned}$$

It follows that

$$(3.4) \quad \tilde{T}(f) = |T|^{\frac{1}{2}} U |T|^{\frac{1}{2}}(f) = \frac{u E(uw)}{E(u^2)} E(uf),$$

for each $f \in L^2(\Sigma)$. Also, we have

$$\begin{aligned} |T^*|^2(f) &= \chi_S w E(u^2) E(wf); \\ |T^*|(f) &= w (E(u^2))^{\frac{1}{2}} (E(w^2))^{-\frac{1}{2}} E(wf) && \text{by Lemma 3.2;} \\ |T^*|^\dagger(f) &= \left(\frac{\chi_S}{E(u^2) (E(w^2))^3} \right)^{\frac{1}{2}} w E(wf) && \text{by (3.2);} \\ U^*(f) &= \left(\frac{\chi_G}{E(u^2) E(w^2)} \right)^{\frac{1}{2}} u E(wf). \end{aligned}$$

Take $r = \chi_G(E(u^2)E(w^2))^{-1/2}$. Then $U^* = M_r M_u E M_w$, and

$$U U^* U = M_w (M_r)^3 M_{E(u^2)} M_{E(w^2)} E M_u = M_r M_w E M_u = U.$$

Thus, U^* is a partial isometry. By (3.1), $N(M_w E M_u) = N(E M_u \sqrt{E(w^2)})$. It follows that

$$\begin{aligned} N(U) &= N(|T|) = N(T); \\ N(U^*) &= N(|T^*|^\dagger) = N(T^\dagger), \end{aligned}$$

and so $T = U|T|$ and $T^\dagger = U^*|T^*|^\dagger$ are the unique polar decompositions.

Theorem 3.16. *Let $T, \tilde{T} \in \text{CR}(L^2(\Sigma))$ with $u, w \geq 0$. Then*

- (a) $T^\dagger = M \frac{u \chi_{\sigma(E(w))}}{E(u^2)E(w^2)} E M_w$.
- (b) $\tilde{T} = M \frac{u E(uw)}{E(u^2)} E M_u$.
- (c) $(\tilde{T})^\dagger = M \frac{u \chi_{\sigma(E(uw))}}{E(u^2)E(uw)} E M_u$.
- (d) $\tilde{T}^\dagger = M \frac{\chi_S w E(uw)}{E(u^2)(E(w^2))^2} E M_w$.
- (e) $\tilde{T}, (\tilde{T})^\dagger$ and \tilde{T}^\dagger are self-adjoint.

Proof. (a) and (b) follows from (3.2) and (3.3).

(c) Take $\nu = \frac{u E(uw)}{E(u^2)}$. Then by (3.3), $\tilde{T} = M_\nu E M_u$. Moreover, by (3.2) we obtain that $(\tilde{T})^\dagger = M \frac{\chi_{\sigma(\nu)}}{E(u^2)E(\nu^2)} M_\nu E M_u$, where $\sigma(\nu) = \sigma(u) \cap \sigma(E(uw))$. Therefore,

$$(\tilde{T})^\dagger(f) = \frac{u \chi_{\sigma(E(uw))}}{E(u^2)E(uw)} E(uf), \quad f \in L^2(\Sigma).$$

(d) Put $\vartheta = \frac{u \chi_G}{E(u^2)E(w^2)}$. Then by (3.2), $T^\dagger = M_\vartheta E M_w$. Hence

$$\tilde{T}^\dagger(f) = \frac{\chi_S w E(uw)}{E(u^2)(E(w^2))^2} E(wf), \quad f \in L^2(\Sigma).$$

(e) It follows from (3.3), (c) and (d). □

Remark 3.17. If we omit the non-negativity hypothesis of u and w in Theorem 3.3, then for every bounded Lambert multiplication $T \in \text{CR}(L^2(\Sigma))$, $\tilde{T}, \tilde{T}^\dagger$ and \tilde{T}^\dagger are always normal operators. Also, by using Theorem 3.3, once again, $\tilde{T}^\dagger = \tilde{T}^\dagger$ if and only if $u = w$.

Let $f \in L^2(\Sigma)$. It is easy to see that

$$\begin{aligned} TT^*(f) &= wE(u^2)E(wf); \\ T^*T(f) &= uE(w^2)E(uf); \\ T^*TTT^*(f) &= uE(w^2)E(u^2)E(uw)E(wf); \\ TT^*T^*T(f) &= wE(u^2)E(w^2)E(uw)E(uf). \end{aligned}$$

So, if $u = w$ or $u, w \in L^0(\mathcal{A})$, then $TT^* = T^*T$. Conversely, if T is normal then

$$(3.5) \quad wE(u^2)E(wf) = uE(w^2)E(uf), \quad f \in L^2(\Sigma).$$

Since \mathcal{A} is sigma-finite, there exists $f_0 \in L^2(\mathcal{A})$ with $\sigma(f_0) = X$. By replacing f_0 by f and then taking the conditional expectation E of both sides of (3.4) gives $E(u^2)(E(w))^2 = (E(u))^2E(w^2)$. Thus, $E(u^2) = (E(u))^2$ and $E(w^2) = (E(w))^2$. But, we know that $E(|f|^2) = |E(f)|^2$ if and only if $f \in L^0(\mathcal{A})$. Consequently, $u, w \in L^0(\mathcal{A})$. Moreover, $T = M_wEM_u$ is binormal if and only if $uE(w) = wE(u)$ on $\sigma(E(uw))$.

We recall that T is an EP operator if and only if $T^\dagger T = TT^\dagger$. Since

$$\begin{aligned} T^\dagger T &= M \frac{\chi_{S \cap G}}{E(u^2)E(w^2)} T^* T; \\ TT^\dagger &= M \frac{\chi_{S \cap G}}{E(u^2)E(w^2)} TT^*, \end{aligned}$$

then T is an EP operator on $L^2(\Sigma)$ if and only if T is a normal operator on $L^2(\Sigma_K)$, where $K = S \cap G$. Thus, we have the following result.

Theorem 3.18. *Let $0 \leq u, w \in L^0(\Sigma)$ with $u \neq w$ and let $T = M_wEM_u \in B(L^2(\Sigma))$. Then the following assertions hold.*

(i) T is normal if and only if $u, w \in L^0(\mathcal{A})$.

(ii) $T \in \text{CR}(L^2(\Sigma))$ is an EP operator on $L^2(\Sigma)$ if and only if $u, w \in L^0(\mathcal{A}_K)$, where $K = S \cap G$.

(iii) T is binormal if and only if $uE(w) = wE(u)$ on $\sigma(E(uw))$.

Now, we determine the lower and upper estimates for the numerical range of T^\dagger . Let $\mu(X) = 1$ and let $T = M_wEM_u \in \text{CR}(L^2(\Sigma))$ with $0 \leq u, w \in \mathcal{D}(E)$. By (3.2) and definition of $\omega(T^\dagger)$ we have

$$\begin{aligned} \omega(T^\dagger) &\geq |\langle T^\dagger 1, 1 \rangle| = \left| \int_X \frac{\chi_{S \cap G}}{E(u^2)E(w^2)} uE(w) d\mu \right| \\ &\geq \int_{S \cap G} \left| \frac{E(u)E(w)}{E(u^2)E(w^2)} \right| d\mu. \end{aligned}$$

On the other hand, since $L^\infty(\Sigma) \cap L^2(\Sigma)$ is dense in $L^2(\Sigma)$, then by the Hölder and conditional Hölder inequality we get that,

$$\begin{aligned} \omega(T^\dagger) &= \sup_{\|f\| \leq 1} |\langle T^\dagger f, f \rangle| \leq \sup_{\|f\| \leq 1} \left| \int_X \frac{\chi_{S \cap G}}{E(u^2)E(w^2)} u E(wf) \bar{f} d\mu \right| \\ &\leq \sup_{\|f\| \leq 1} \int_X \left| \frac{\chi_{S \cap G}}{E(u^2)E(w^2)} (E(u^2))^{\frac{1}{2}} (E(w^2))^{\frac{1}{2}} E(|f|^2) \right| d\mu \\ &\leq \sup_{\|f\| \leq 1} \int_{S \cap G} \frac{1}{(E(u^2))^{\frac{1}{2}} (E(w^2))^{\frac{1}{2}}} E(|f|^2) d\mu \\ &\leq \int_{S \cap G} \frac{d\mu}{(E(u^2))^{\frac{1}{2}} (E(w^2))^{\frac{1}{2}}}. \end{aligned}$$

Consequently, we have the following theorem.

Theorem 3.19. *Let $\mu(X) = 1$ and let $T = M_w E M_u \in \text{CR}(L^2(\Sigma))$ with $0 \leq u, w \in \mathcal{D}(E)$. Then*

$$\int_{S \cap G} \frac{E(u)E(w)}{E(u^2)E(w^2)} d\mu \leq \omega(T^\dagger) \leq \int_{S \cap G} \frac{d\mu}{\sqrt{E(u^2)E(w^2)}},$$

where $S = \sigma(E(u))$, $G = \sigma(E(w))$.

Example 3.20. Let $X = [-\frac{1}{2}, \frac{1}{2}]$, $d\mu = dx$, Σ be the Lebesgue sets, and let $\mathcal{A} \subseteq \Sigma$ be the σ -algebra generated by the symmetric sets about the origin. Let $0 < a \leq \frac{1}{2}$ and $f \in L^2(\Sigma)$. Then

$$\begin{aligned} \int_{-a}^a E(f)(x) dx &= \int_{-a}^a f(x) dx \\ &= \int_{-a}^a \left\{ \frac{f(x) + f(-x)}{2} + \frac{f(x) - f(-x)}{2} \right\} dx = \int_{-a}^a \frac{f(x) + f(-x)}{2} dx. \end{aligned}$$

Thus, $E(f)(x) = \frac{f(x) + f(-x)}{2}$. Put $u(x) = x + 2$, $w(x) = x + 3$ and $T = M_w E M_u$. Then $E(u) = 2$, $E(w) = 3$, $E(u^2) = x^2 + 4$ and $E(w^2) = x^2 + 9$. Now, by Proposition 3.1(i) we get that

$$\|T\| = \|\sqrt{(x^2 + 4)(x^2 + 9)}\|_\infty = \frac{\sqrt{629}}{4} = 6.2699.$$

Moreover, since $u\sqrt{E(w^2)} = (x + 2)\sqrt{x^2 + 9} \geq \frac{9}{2}$, then by Proposition 3.1(ii),

$T \in \text{CR}(L^2(\Sigma))$. Also, it is easy to check that

$$\begin{aligned} \int_{[-\frac{1}{2}, \frac{1}{2}]} \frac{E(u)E(w)}{E(u^2)E(w^2)} d\mu &= \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{6dx}{(x^2+4)(x^2+9)} = 0.1618; \\ \int_{[-\frac{1}{2}, \frac{1}{2}]} \frac{d\mu}{\sqrt{E(u^2)E(w^2)}} &= \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{dx}{\sqrt{(x^2+4)(x^2+9)}} = 0.1642; \\ \|T^\dagger\| &= \left\| \frac{1}{\sqrt{E(u^2)E(w^2)}} \right\|_\infty = \frac{1}{6} = 1.666; \\ \|\tilde{T}\| &= \|E(uw)\|_\infty = \frac{25}{4} = 6.250. \end{aligned}$$

Thus, $\|\tilde{T}\| \leq \|T\|$, $\|T^\dagger\| \geq 1/\|T\|$ and by Theorem 3.6 we obtain

$$0.1618 \leq \omega(T^\dagger) \leq 0.1642 \leq 1.666 = \|T^\dagger\|.$$

Acknowledgements

The authors would like to thank the referee for very helpful comments and valuable suggestions.

REFERENCES

1. A. Aluthge, On p -hyponormal operators for $0 < p < 1$, Integral Equations Operator Theory **13** (1990), 307-315.
2. A. Ben-Israel and T. N. E. Greville, Generalized inverses: Theory and Applications, Springer-Verlag, New York, 2003.
3. J. Benda and S. Sherman, Monotone and convex operator functions, Trans. Amer. Math. Soc. **79** (1955), 58-71.
4. A. Brown, C. K. Fong and D. W. Hadwin, Parts of operators on Hilbert space, Illinois J. Math. **22** (1978), 306-314.
5. Y. Estaremi, On a class of operators with normal Aluthge transformations, Filomat **29** (2015), 969-975.
6. Y. Estaremi, Some classes of weighted conditional type operators and their spectra, Positivity **19** (2015), 83-93.
7. M. Cho, R. E. Curto and T. Huruya, n -tuples of operators satisfying $\sigma_T(AB) = \sigma_T(BA)$, Linear Algebra Appl. **341** (2002), 291-298.
8. D. S. Djordjević, Products of EP operators on Hilbert spaces, Proc. Amer. Math. Soc. **129** (2001), 1727-1731.
9. D. S. Djordjević, Characterizations of normal, hyponormal and EP operators, J. Math. Anal. Appl. **329** (2007), 1181-1190.
10. D. S. Djordjević and N. C. Dinčić, Reverse order law for the Moore-Penrose inverse, J. Math. Anal. Appl. **361** (2010), 252-261.

11. T. Furuta, On the polar decomposition of an operator, *Acta Sci. Math. (Szeged)* **46** (1983), 261-268.
12. T. Furuta, *Invitation to linear operators*, Taylor & Francis, Ltd. London, 2001.
13. F. Gesztesy, M. Malamud, M. Mitrea and S. Naboko, Generalized polar decompositions for closed operators in Hilbert spaces and some applications, *Integral Equations Operator Theory* **64** (2009), 83-113.
14. J. Herron, Weighted conditional expectation operators, *Oper. Matrices* **5** (2011), 107-118.
15. M. R. Jabbarzadeh, A conditional expectation type operator on L^p spaces, *Oper. Matrices* **4** (2010), 445-453.
16. S. Jung, Y. Kim and E. Ko, Composition operators for which $C_\phi^*C_\phi$ and $C_\phi + C_\phi^*$ commute, *Complex Var. Elliptic Equ.* **59** (2014), 1608-1625.
17. F. Kimura, Analysis of non-normal operators via Aluthge transformation, *Integral Equations Operator Theory* **50** (2004), 375-384.
18. M. M. Rao, *Conditional measure and applications*, Marcel Dekker, New York, 1993.
19. T. Yamazaki, Parallelisms between Aluthge transformation and powers of operators, *Acta Sci. Math. (Szeged)* **67**(2001), 809-820.
20. T. Yamazaki, On numerical range of the Aluthge transformation, *Linear Algebra Appl.* **341** (2002), 111-117.
21. T. Yamazaki, Characterizations of $\log A \geq \log B$ and normaloid operators via Heinz inequality, *Integral Equations Operator Theory* **43** (2002), 237-247.
22. T. Yamazaki, On the polar decomposition of the Aluthge transformation and related results, *J. Operator Theory* **51** (2004), 303-319.

Mohammad Reza Jabbarzadeh

University of Tabriz
Faculty of Mathematical Sciences
Tabriz, Iran,
E-mail: *mjabbar@tabrizu.ac.ir*

(Received 26.10.2016)

(Revised 23.02.2018)

Hossein Emamalipour

University of Tabriz
Faculty of Mathematical Sciences
Tabriz, Iran,
E-mail: *h_emamali@tabrizu.ac.ir*

Morteza Sohrabi Chegeni

Lorestan University
Department of Mathematics
Khorramabad, Iran,
E-mail: *mortezasohrabi021@gmail.com*