

WEIGHTED P–PARTITIONS ENUMERATOR

Marko Pešović and Tanja Stojadinović

To an extended generalized permutohedron we associate the weighted integer points enumerator, whose principal specialization is the f -polynomial. In the case of poset cones it refines Gessel’s P-partitions enumerator. We show that this enumerator is a quasisymmetric function obtained by universal morphism from the Hopf algebra of posets.

1. INTRODUCTION

In the seminal paper of Aguiar, Bergeron and Sottile [2] the notion of combinatorial Hopf algebra was introduced. They explained the ubiquity of quasisymmetric functions as generating functions in enumerative combinatorics. More recently a geometric meaning of quasisymmetric enumerators is attributed. It is based on a class of convex polytopes called generalized permutohedra. The integer points enumerator associated to a generalized permutohedron is a quasisymmetric function. It was defined, and studied in the case of matroid base polytopes, by Billera, Jia and Reiner in [3], and in the case of nestohedra by Grujić in [6]. More subtle generalization which takes into account the face structure of a generalized permutohedron is introduced and studied in [8]. In this paper we consider the extended generalized permutohedra and the special case of poset cones. We prove that the integer points enumerator associated to a poset cone coincides with the universal morphism from the Hopf algebra of posets to quasisymmetric functions. The specialization for $q = 0$ is the Gessel enumerator of P-partitions which attributes the geometric meaning to this classical function.

2020 Mathematics Subject Classification. 16T05, 52B05

Keywords and Phrases. Generalized permutohedron, Quasisymmetric function, Poset, Combinatorial Hopf algebra.

In section 2 we review necessary facts about quasisymmetric functions and \mathcal{P} -partitions enumerators. In section 3 we introduce the integer points enumerator $F_q(P)$ for an extended generalized permutohedron P , which is a weighted quasisymmetric function, in the same manner as in the case of generalized permutohedra provided in [8]. The parameter q reflects the rank function of the face lattice $L(P)$. To a poset \mathcal{P} is associated the poset cone $C(\mathcal{P})$, which is an extended generalized permutohedron and our construction produces a weighted quasisymmetric function $F_q(C(\mathcal{P}))$. In section 4 we prove Theorem 21, the first main result of the paper, which states that the weighted quasisymmetric function $F_q(C(\mathcal{P}))$, constructed geometrically, has an algebraic meaning as the universal morphism from a certain combinatorial Hopf algebra of posets \mathcal{P} to the Hopf algebra $QSym$ of quasisymmetric functions. This result is analogous to the previous results for simple graphs [7], matroids and building sets [8], and spreads their validity to the case of extended generalized permutohedra. The main theorem is followed by various examples, and statements about behavior of the enumerator of the opposite poset and under the action of the antipode. In Theorem 37 in section 5, it is shown that for a well labelled poset \mathcal{P} and $q = 0$ our enumerator specializes to the classical Gessel's \mathcal{P} -partitions enumerator. We also provide an example of posets with the same \mathcal{P} -partitions enumerators but which are distinguished by corresponding weighted quasisymmetric enumerators.

2. QUASISYMMETRIC FUNCTIONS

A *composition* α of a positive integer n , $\alpha \models n$, is an ordered list $(\alpha_1, \dots, \alpha_k)$ of positive integers such that $\alpha_1 + \dots + \alpha_k = n$. The *monomial quasisymmetric function* M_α indexed by the composition α is an element of the commutative algebra of formal power series in the countable ordered set of variables $\mathbf{x} = (x_1 < x_2 < x_3 < \dots)$ defined by

$$M_\alpha = \sum_{i_1 < \dots < i_k} x_{i_1}^{\alpha_1} \cdots x_{i_k}^{\alpha_k}.$$

The *algebra of quasisymmetric functions* $QSym$, spanned by M_α when α runs over all compositions, is a subalgebra of the algebra of formal power series. The algebra $QSym$ is a graded, connected Hopf algebra (see [5], Proposition 5.1.8). The homogeneous component $QSym_n$ is spanned by $\{M_\alpha\}_{\alpha \models n}$. Let $\zeta_Q : QSym \rightarrow \mathbf{k}$ be a linear multiplicative functional defined on the monomial basis by

$$\zeta_Q(M_\alpha) = \begin{cases} 1, & \text{if } \alpha = (n) \text{ for } n \in \mathbb{N}, \\ 0, & \text{otherwise.} \end{cases}$$

The Hopf algebra $QSym$ equipped with the character ζ_Q is the terminal object in the category of combinatorial Hopf algebras. For a definition of the combinatorial Hopf algebra see [2].

Theorem 1 ([2], Theorem 4.1). *For a combinatorial Hopf algebra (\mathcal{H}, ζ) there is a unique morphism of graded Hopf algebras $\Psi : H \rightarrow \mathcal{QSym}$ such that*

$$\zeta_{\mathcal{Q}} \circ \Psi = \zeta.$$

For a homogeneous element h of degree n the coefficients $\zeta_{\alpha}(h)$, $\alpha = (\alpha_1, \dots, \alpha_k) \models n$ of $\Psi(h)$ in the monomial basis are given by

$$\zeta_{\alpha}(h) = \zeta^{\otimes k} \circ (p_{\alpha_1} \otimes \dots \otimes p_{\alpha_k}) \circ \Delta^{k-1}(h),$$

where p_i is the projection of \mathcal{H} on the i -th homogeneous component \mathcal{H}_i and Δ^{k-1} is the $(k-1)$ -fold coproduct map of \mathcal{H} .

For $F \in \mathcal{QSym}$ and $m \in \mathbb{N}$, let \mathbf{ps}^1 denotes the *principal specialization*

$$\mathbf{ps}^1(F)(m) = F(\underbrace{1, \dots, 1}_m, 0, 0 \dots).$$

Note that $\mathbf{ps}^1(F)(m)$ is a polynomial on a variable m and

$$\mathbf{ps}^1(M_{\alpha})(m) = \binom{m}{k(\alpha)},$$

where $k(\alpha)$ is the number of parts of $\alpha = (\alpha_1, \dots, \alpha_k)$. Specially, for $m = -1$ we have

$$\mathbf{ps}^1(M_{\alpha})(-1) = \binom{-1}{k(\alpha)} = (-1)^{k(\alpha)}.$$

For a composition $\alpha = (\alpha_1, \dots, \alpha_k) \models n$ let $D(\alpha) \subseteq [n-1]$ be a subset defined by $D(\alpha) = \{\alpha_1, \alpha_1 + \alpha_2, \dots, \alpha_1 + \alpha_2 + \dots + \alpha_{k-1}\}$. We say that $\alpha \models n$ *refines* $\beta \models n$, and write $\beta \preceq \alpha$, if $D(\beta) \subseteq D(\alpha)$.

Another important basis of \mathcal{QSym} is the basis of *fundamental quasisymmetric functions* defined by

$$L_{\alpha} = \sum_{\alpha \preceq \beta} M_{\beta}.$$

2.1 Quasisymmetric enumerator of P-partitions

A *labelled poset* P is a poset on some finite subset of positive integers. A *P-partition* is a function $f : P \rightarrow \mathbb{N}$ such that

- $i <_P j$ and $i <_{\mathbb{Z}} j$ implies $f(i) \leq f(j)$,
- $i <_P j$ and $i >_{\mathbb{Z}} j$ implies $f(i) < f(j)$.

Definition 2. A poset \mathbf{P} is a *well labelled poset* if $i <_{\mathbf{P}} j$ implies $i >_{\mathbb{Z}} j$. In that case, a \mathbf{P} -partition is a function $f : \mathbf{P} \rightarrow \mathbb{N}$ such that

$$i <_{\mathbf{P}} j \quad \text{implies} \quad f(i) < f(j).$$

Denote by $\mathcal{A}(\mathbf{P})$ the set of all \mathbf{P} -partitions. For a poset \mathbf{P} on $[n]$, define the *enumerator of \mathbf{P} -partitions* by

$$F_{\mathbf{P}}(\mathbf{x}) = \sum_{f \in \mathcal{A}(\mathbf{P})} x_{f(1)} x_{f(2)} \cdots x_{f(n)}.$$

Proposition 3 ([5], Proposition 5.2.10). *For a totally ordered labelled poset $\mathbf{P} = \{i_1 <_{\mathbf{P}} i_2 <_{\mathbf{P}} \cdots <_{\mathbf{P}} i_n\}$ the enumerator of \mathbf{P} -partitions is equal to the fundamental quasisymmetric function*

$$F_{\mathbf{P}}(\mathbf{x}) = L_{\alpha(\mathbf{P})},$$

where $\alpha(\mathbf{P}) \models n$ is a composition such that $D(\alpha(\mathbf{P})) = \{j : i_j >_{\mathbb{Z}} i_{j+1}\}$.

Theorem 4 ([5], Theorem 5.2.11). *For a labelled poset \mathbf{P} ,*

$$F_{\mathbf{P}}(\mathbf{x}) = \sum_{\mathbf{l} \in \mathcal{L}(\mathbf{P})} F_{\mathbf{l}}(\mathbf{x}),$$

where the sum is over the set $\mathcal{L}(\mathbf{P})$ of all linear extensions \mathbf{l} of \mathbf{P} .

Example 5. Figure 1 presents two posets with their enumerators of \mathbf{P} -partitions. Note that the poset \mathbf{P}_1 is not well labelled, while the poset \mathbf{P}_2 is.




Figure 1: $F_{\mathbf{P}_1}(\mathbf{x}) = L_{(1,3)} + L_{(2,2)}$ and $F_{\mathbf{P}_2}(\mathbf{x}) = L_{(2,1,1)} + L_{(1,1,1,1)}$

3. POSET CONES

To a finite poset \mathbf{P} we can associate its polyhedral model called the poset cone. It belongs to a wider class of polyhedra called extended generalized permutohedra.

A standard $(n - 1)$ -dimensional permutohedron Pe^{n-1} is the convex hull of the orbit of a point $(a_1, a_2, \dots, a_n) \in \mathbb{R}^n$ with increasing coordinates $a_1 < a_2 < \dots < a_n$

$$Pe^{n-1} = \text{conv}\{(a_{\sigma(1)}, a_{\sigma(2)}, \dots, a_{\sigma(n)}) : \sigma \in \mathfrak{S}_n\},$$

where \mathfrak{S}_n is the permutation group of $[n]$.

Proposition 6 ([4], Proposition 1.3, [10], Proposition 2.6). *The d -dimensional faces of Pe^{n-1} are in one-to-one correspondence with order compositions $\mathcal{C} = C_1|C_2|\dots|C_{n-d}$ of $[n]$. The affine hull of the face corresponding to the order composition \mathcal{C} is given by the $n - d$ linear equations*

$$\sum_{i \in C_1 \cup C_2 \cup \dots \cup C_k} x_i = a_1 + a_2 + \dots + a_{|C_1 \cup C_2 \cup \dots \cup C_k|},$$

for $1 \leq k \leq n - d$.

An order composition $\mathcal{C} = C_1|C_2|\dots|C_{n-d}$ defines the *flag*, or *nested set*, \mathcal{F} of subsets

$$\mathcal{F} : \emptyset = F_0 \subset F_1 \subset \dots \subset F_{n-d-1} \subset F_{n-d} = [n],$$

of length $|\mathcal{F}| = n - d$, where $F_i = C_1 \cup C_2 \cup \dots \cup C_i$, for $1 \leq i \leq n - d$. There is an obvious order reversing one-to-one correspondence between the face lattice of the permutohedron $L(Pe^{n-1})$ and the lattice of flags of subsets of the set $[n]$. Using this correspondence we will label faces of the standard permutohedron Pe^{n-1} by flags of subsets of $[n]$. We have

$$\dim(\mathcal{F}) = n - |\mathcal{F}|.$$




Figure 2: The permutohedron Pe^2 and the corresponding braid arrangement

The *normal fan* $\mathcal{N}(P)$ of a polyhedron P is the collection of normal cones to faces of P . Recall that the *normal cone* to a face G is given by

$$C_G := \{\omega \in \mathbb{R}^n : G \subseteq \{x \in P : \omega^*(x) = \max\{\omega^*(y) : y \in P\}\}\}.$$

The normal fan $\mathcal{N}(Pe^{n-1})$ of the standard permutohedron Pe^{n-1} is the fan of the *braid arrangement*, given by hyperplanes $\{x_i = x_j\}_{1 \leq i < j \leq n}$, in \mathbb{R}^n . The cones of the braid arrangement fan are called *braid cones*. The braid cone $C_{\mathcal{F}}$ at the face \mathcal{F} is determined by

- $x_p = x_q$ if $p, q \in F_{i+1} \setminus F_i$, for some $0 \leq i \leq |\mathcal{F}| - 1$,
- $x_p \leq x_q$ if $p \in F_i \setminus F_{i-1}$ and $q \in F_{i+1} \setminus F_i$, for some $1 \leq i \leq |\mathcal{F}| - 1$.

We have $\dim(C_{\mathcal{F}}) = |\mathcal{F}|$. The fan \mathcal{N}_1 is a *refinement* of \mathcal{N}_2 (or \mathcal{N}_2 is a *coarsening* of \mathcal{N}_1) if every cone of \mathcal{N}_1 is contained in a cone in \mathcal{N}_2 (or if every cone in \mathcal{N}_2 is a union of cones of \mathcal{N}_1).

Definition 7. A convex polytope Q is an $(n-1)$ -dimensional *generalized permutohedron* if the braid arrangement fan $\mathcal{N}(Pe^{n-1})$ refines the normal fan $\mathcal{N}(Q)$.




Figure 3: Generalized permutohedron

For a generalized permutohedron Q there is a map $\pi_Q : L(Pe^{n-1}) \rightarrow L(Q)$ between face lattices, determined by $\pi_Q(\mathcal{F}) = G$ if and only if the relative interior of the braid cone $C_{\mathcal{F}}^\circ$ is contained in the relative interior C_G° of the normal cone C_G at the face $G \in L(Q)$. We say that the flag \mathcal{F} is *normal* to the face G . Denote by $F(G) = \{\mathcal{F} : C_{\mathcal{F}}^\circ \subseteq C_G^\circ\}$ the set of normal flags to a face G . By Proposition 2.3 in [8], we have

$$(1) \quad \sum_{\mathcal{F} \in F(G)} (-1)^{|\mathcal{F}|} = (-1)^{n-1-\dim(G)}.$$

Definition 8. An *extended generalized permutohedron* P is a polyhedron whose normal fan $\mathcal{N}(P)$ is a coarsening of a subfan of the braid arrangement fan.

An example of an extended generalized permutohedron is given on Figure 4. We are concerned on the class of extended generalized permutohedra associated to finite posets. We follow the description given in [1].

Definition 9 ([1], Proposition 15.1). The *poset cone* of a poset P on $[n]$ is an extended generalized permutohedron given by

$$C(P) = \text{cone}\{e_i - e_j : i <_P j\},$$




Figure 4: The extended generalized permutohedron

where e_1, e_2, \dots, e_n are the standard basis vectors in \mathbb{R}^n .

Proposition 10 ([1], Proposition 15.1). *The generating rays of $C(P)$ are determined by the vectors $e_i - e_j$ corresponding to the cover relations $i <_P j$ in P .*

A subset $S \subseteq [n]$ is called an *ideal* of P , denoted by $S \triangleleft P$, if no element of $[n] \setminus S$ is less than an element of S . It is shown in [1] that the poset cone is described by

$$C(P) = \left\{ (x_1, \dots, x_n) : \sum_{i=1}^n x_i = 0 \text{ and } \sum_{s \in S} x_s \geq 0 \text{ for all } S \triangleleft P \right\}.$$

Therefore

$$(2) \quad \dim(C(P)) = n - c(P),$$

where $c(P)$ is the number of connected components of the poset P . Recall that a poset is said to be *connected* if for all pair $i_1, i_2 \in [n]$ there is a finite sequence $i_j = j_1, j_2, \dots, j_m = i_2 \in [n]$ such that for all i $j_k \leq_P j_{k+1}$ or $j_k \geq_P j_{k+1}$. A *connected component* of P is a maximal connected subposet.

Let $C : i_1, i_2, \dots, i_n$ be a cyclic sequence of elements of P , the poset on the set $[n]$, where every consecutive pair is comparable in P . We say that C is a *circuit* of P . Circuits consist of *up-edges* $i_j <_P i_{j+1}$ and *down-edges* $i_j >_P i_{j+1}$.

Definition 11. A subposet Q of a poset P on $[n]$ is *positive* if for every circuit C all down-edges of C are in Q if and only if all up-edges of C are in Q . Let $\text{Pos}(P)$ be the set of all positive subposets of P .

The faces of the poset cone $C(P)$ are characterized by the following lemma.

Lemma 12 ([1], Lemma 15.3). *Let P be a poset on $[n]$. The faces of the poset cone $C(P) \subseteq \mathbb{R}^n$ are precisely the poset cones $C(Q)$ as Q ranges over positive subposets of P .*

The following proposition describes the normal fan $\mathcal{N}(C(\mathbf{P}))$ of the poset cone $C(\mathbf{P})$. It shows that it coincides with the braid cone $\sigma_{\mathbf{P}}$ considered in [11, Proposition 3.5], so we can call it the normal cone of the poset cone.

Proposition 13. *A vector $\omega = (\omega_1, \dots, \omega_n) \in \mathbb{Z}_+^n$ lies in $\mathcal{N}(C(\mathbf{P}))$ if and only if $\omega_i \leq \omega_j$ for any $i \leq_{\mathbf{P}} j$.*

Proof. A vector $\omega \in \mathbb{Z}_+^n$ lies in $\mathcal{N}(C(\mathbf{P}))$ if and only if ω^* is maximized over $C(\mathbf{P})$ at some of its face. The weight function ω^* on the ray generated by $e_i - e_j$ is given by

$$\omega^*(t(e_i - e_j)) = \langle \omega, t(e_i - e_j) \rangle = t(\omega_i - \omega_j), t \geq 0.$$

Consequently, we have:

1. If $\omega_i > \omega_j$, then $t(\omega_i - \omega_j) \geq 0$ for $t \geq 0$ and maximum of ω^* is not achieved along the ray generated by $e_i - e_j$.
2. If $\omega_i < \omega_j$, then $t(\omega_i - \omega_j) \leq 0$ for $t \geq 0$ and maximum along the ray generated by $e_i - e_j$ is achieved at the vertex of $C(\mathbf{P})$.
3. If $\omega_i = \omega_j$, then $t(\omega_i - \omega_j) = 0$ for $t \geq 0$ and ω^* is maximized along the ray generated by $e_i - e_j$.

□

Corollary 14. *A flag \mathcal{F} is normal to the face $C(\mathbf{Q})$ for some positive subposet $\mathbf{Q} \in \text{Pos}(\mathbf{P})$, i.e. $\mathcal{F} \in \mathbf{F}(C(\mathbf{Q}))$ if and only if for each $\omega \in C_{\mathcal{F}}^{\circ}$,*

1. $\omega_i = \omega_j$ for all $i \leq_{\mathbf{Q}} j$,
2. $\omega_i < \omega_j$ for all i, j which are incomparable in \mathbf{Q} and $i \leq_{\mathbf{P}} j$.

The following example illustrates the concepts introduced in connection with poset cones.

Example 15. Let \mathbf{P} be a poset on $[4]$ defined by covering relations $1, 2 < 3, 4$. The generated rays of the poset cone $C(\mathbf{P})$ are determined by vectors $e_1 - e_3, e_1 - e_4, e_2 - e_3, e_2 - e_4$. The normal fan $\mathcal{N}(C(\mathbf{P}))$ is described by inequalities $x_1, x_2 \leq x_3, x_4$, see Figure 5.

The list of circuits of \mathbf{P} is the following

$$\begin{aligned} &1 < 4 > 2 < 3, 1 < 3 > 2 < 4, 4 > 2 < 3 > 1, 3 > 2 < 4 > 1, \\ &2 < 3 > 1 < 4, 2 < 4 > 1 < 3, 3 > 1 < 4 > 2, 4 > 1 < 3 > 2. \end{aligned}$$

Its facets are determined by positive subposets given by

$$1 < 3, 4; 2 < 3, 4; 1, 2 < 3; 1, 2 < 4.$$




Figure 5: Poset cone and its normal fan

3.2 Quasisymmetric enumerator $F_q(P)$

To an extended generalized permutohedron P we associate the weighted enumerator $F_q(P)$ of positive integer points in its normal fan. A vector $\omega = (\omega_1, \omega_2, \dots, \omega_n) \in \mathbb{Z}_+^n$ defines the *weight function* $\omega^* : \mathbb{R}^n \rightarrow \mathbb{R}$ by $\omega^*(x) := \langle \omega, x \rangle$ where $\langle \cdot, \cdot \rangle$ is the standard scalar product in \mathbb{R}^n . The weight function ω^* on Pe^{n-1} is maximized along a unique face \mathcal{F}_ω of Pe^{n-1} determined by the condition that the vector ω lies in the relative interior of its normal cone $\omega \in C_{\mathcal{F}_\omega}^\circ$. The flag \mathcal{F}_ω satisfies the following conditions

- ω is constant on $F_i \setminus F_{i-1}$, for $1 \leq i \leq k$,
- $\omega|_{F_i \setminus F_{i-1}} < \omega|_{F_{i+1} \setminus F_i}$, for $1 \leq i \leq k-1$,

where k is the length of \mathcal{F}_ω . Let $M_{\mathcal{F}}$ be the *enumerator* of positive integer vectors $\omega \in \mathbb{Z}_+^n$ in relative interior of the corresponding braid cone $C_{\mathcal{F}}$

$$(3) \quad M_{\mathcal{F}} = \sum_{\omega \in \mathbb{Z}_+^n \cap C_{\mathcal{F}}^\circ} x_{\omega_1} x_{\omega_2} \cdots x_{\omega_n}.$$

Note that $\omega \in C_{\mathcal{F}}^\circ$ if and only if $\mathcal{F}_\omega = \mathcal{F}$. The enumerator $M_{\mathcal{F}}$ is a monomial quasisymmetric function indexed by composition

$$\text{type}(\mathcal{F}) = (|F_1 \setminus F_0|, |F_2 \setminus F_1|, \dots, |F_k \setminus F_{k-1}|).$$

A *weight function* ω^* on an extended generalized permutohedron P is maximized along a unique face G_ω of P which is determined by the condition $\omega \in C_{G_\omega}^\circ$. For a face G of P we define a *quasisymmetric enumerator* $F_q(G)$ by

$$F_q(G) = q^{\dim(G)} \sum_{\omega \in \mathbb{Z}_+^n \cap C_G^\circ} x_{\omega_1} x_{\omega_2} \cdots x_{\omega_n}.$$

By (3) it follows immediately that

$$F_q(G) = q^{\dim(G)} \sum_{\mathcal{F} \in \mathcal{F}(G)} M_{\mathcal{F}}.$$

Definition 16. For an extended generalized permutohedron P with the face lattice $L(P)$ let

$$F_q(P) = \sum_{G \in L(P)} F_q(G).$$

The following proposition describes the f -polynomial $f(P, q) := f_0 + f_1q + f_2q^2 + \cdots + f_{n-1}q^{n-1}$ of an extended generalized permutohedron P in terms of the weighted quasisymmetric enumerator $F_q(P)$.

Proposition 17. *The f -polynomial of an $(n-1)$ -dimensional extended generalized permutohedron P is determined by the principal specialization*

$$f(P, q) = (-1)^{n-1} \mathbf{ps}^1(F_{-q}(P))(-1).$$

Proof. It follows from (1) that

$$\mathbf{ps}^1(F_{-q}(G))(-1) = (-1)^{n-1} q^{\dim(G)}.$$

Therefore

$$\mathbf{ps}^1(F_{-q}(P))(-1) = (-1)^{n-1} \sum_{G \in L(P)} q^{\dim(G)}.$$

□

Quasisymmetric enumerator for a poset cone

By Definition 16 and Lemma 12, it follows that the weighted quasisymmetric enumerator $F_q(C(P))$ for a poset P can be expressed as

$$(4) \quad F_q(C(P)) = \sum_{Q \in \text{Pos}(P)} \sum_{\mathcal{F} \in \mathcal{F}(C(Q))} q^{\dim(C(Q))} M_{\mathcal{F}}.$$

Example 18. Direct calculation of quasisymmetric enumerator for a poset cone in Example 15 gives

$$\begin{aligned} F_q(C(P)) &= q^3 M_{(4)} + 2q^2 (M_{(1,3)} + M_{(3,1)}) + 4q M_{(1,2,1)} \\ &\quad + M_{(2,2)} + 2M_{(1,1,2)} + 2M_{(2,1,1)} + 4M_{(1,1,1,1)}. \end{aligned}$$

The corresponding f -polynomial is $f(C(P), q) = q^3 + 4q^2 + 4q + 1$.

4. HOPF ALGEBRA \mathcal{P}

The set of isomorphism classes of finite posets linearly generates the \mathbf{k} -vector space \mathcal{P}

$$\mathcal{P} = \bigoplus_{n \geq 0} \mathcal{P}_n,$$

where \mathcal{P}_n is the homogeneous component of degree n spanned by isomorphism classes of posets $[P]$ on $[n]$. The space \mathcal{P} is a graded, commutative and non-cocommutative Hopf algebra (see [2], Example 2.3) with the *multiplication* and the *comultiplication* given by

$$[P_1] \cdot [P_2] = [P_1 \sqcup P_2],$$

$$\Delta([P]) = \sum_{S \triangleleft P} [P|_S] \otimes [P|_{[n]\setminus S}],$$

where $P_1 \sqcup P_2$ is the disjoint union of posets, and $P|_S$ is the restriction of the poset to the subset $S \subset [n]$.

Definition 19. A flag $\mathcal{F} : \emptyset = F_0 \subset F_1 \subset \dots \subset F_k = [n]$ is a *flag of ideals* of a poset P , denoted by $\mathcal{F} \triangleleft P$, if $F_i \triangleleft P$, for all $0 < i \leq k$. Let

$$\mathfrak{F}(P) = \{\mathcal{F} : \mathcal{F} \triangleleft P\}.$$

There is a map $\mathfrak{F}(P) \rightarrow \mathcal{P}_n$ given by

$$\mathcal{F} \mapsto P/\mathcal{F} = \prod_{i=1}^k P|_{F_i \setminus F_{i-1}}.$$

Example 20. Let $\mathcal{F} : \emptyset \subset \{1, 2, 5\} \subset \{1, 2, 4, 5, 7\} \subset [8]$.




Figure 6: Posets P and P/\mathcal{F}

A poset P/\mathcal{F} is a weak subposet of P , i.e. if $i \leq j$ in P/\mathcal{F} then $i \leq j$ in P . Let $P/\mathcal{F} = P_1 \sqcup P_2 \sqcup \dots \sqcup P_m$ be the decomposition into connected components. We say that P_u and P_v are *incomparable* if elements from P_u and P_v are mutually incomparable in a poset P . Otherwise, P_u is *smaller* than P_v if no element of P_v is less than an element of P_u .

4.3 Quasisymmetric enumerator $F_q(C(\mathbf{P}))$

We extend the basic field \mathbf{k} into the field of rational functions $\mathbf{k}(q)$ and define the character $\zeta_q : \mathcal{P} \rightarrow \mathbf{k}(q)$ with

$$\zeta_q([\mathbf{P}]) = q^{\text{rk}(\mathbf{P})} \quad \text{where} \quad \text{rk}(\mathbf{P}) = n - c(\mathbf{P}).$$

Let $\Psi_q : (\mathcal{P}, \zeta_q) \rightarrow (\mathcal{QSym}, \zeta_{\mathcal{Q}})$ be the unique morphism of combinatorial Hopf algebras over $\mathbf{k}(q)$, given by Theorem 1 with

$$(5) \quad \Psi_q([\mathbf{P}]) = \sum_{\alpha \models n} (\zeta_q)_\alpha(\mathbf{P}) M_\alpha.$$

The coefficient corresponding to a composition $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k) \models n$ is determined by

$$(\zeta_q)_\alpha(\mathbf{P}) = \sum_{\substack{\mathcal{F} \in \mathfrak{F}(\mathbf{P}) \\ \text{type}(\mathcal{F}) = \alpha}} \prod_{j=1}^k q^{\text{rk}(\mathbf{P}|_{F_j \setminus F_{j-1}})} = \sum_{\substack{\mathcal{F} \in \mathfrak{F}(\mathbf{P}) \\ \text{type}(\mathcal{F}) = \alpha}} q^{\text{rk}_{\mathbf{P}}(\mathcal{F})},$$

where

$$\text{rk}_{\mathbf{P}}(\mathcal{F}) = \sum_{j=1}^k \text{rk}(\mathbf{P}|_{F_j \setminus F_{j-1}}) = n - \sum_{j=1}^k c(\mathbf{P}|_{F_j \setminus F_{j-1}}).$$

Thus, the equation (5) can be expressed as

$$\Psi_q([\mathbf{P}]) = \sum_{\mathcal{F} \in \mathfrak{F}(\mathbf{P})} q^{\text{rk}_{\mathbf{P}}(\mathcal{F})} M_{\mathcal{F}}.$$

To a poset \mathbf{P} are associated two weighted quasisymmetric functions, $F_q(C(\mathbf{P}))$ with a geometric meaning based on the combinatorics of the poset cone $C(\mathbf{P})$ and $\Psi_q([\mathbf{P}])$ with an algebraic meaning based on the Hopf algebra structure on finite posets. The following theorem shows that these two functions are equal.

Theorem 21. *For a poset \mathbf{P} the quasisymmetric enumerator function $F_q(C(\mathbf{P}))$ associated to a poset cone $C(\mathbf{P})$ coincides with the value of the universal morphism at $[\mathbf{P}]$ from the combinatorial Hopf algebra of posets \mathcal{P} to \mathcal{QSym}*

$$F_q(C(\mathbf{P})) = \Psi_q([\mathbf{P}]).$$

Proof. The weighted enumerator $F_q(C(\mathbf{P}))$ is described by (4). We need to show that

$$\mathfrak{F}(\mathbf{P}) = \{\mathcal{F} : \mathcal{F} \in \mathbf{F}(C(\mathbf{Q})) \text{ for some } \mathbf{Q} \in \text{Pos}(\mathbf{P})\},$$

and $\text{rk}_{\mathbf{P}}(\mathcal{F}) = \dim(C(\mathbf{Q}))$, for $\mathcal{F} \in \mathbf{F}(C(\mathbf{Q}))$. Let \mathbf{P} be a poset on the set $[n]$.

\supseteq : Suppose $\mathcal{F} \in \mathbf{F}(C(\mathbf{Q}))$ for some $\mathbf{Q} \in \text{Pos}(\mathbf{P})$. If $\mathbf{Q} = \mathbf{Q}_1 \sqcup \dots \sqcup \mathbf{Q}_m$ is the decomposition into connected components, by (2) we have $\dim(C(\mathbf{Q})) = n - m$. For $\omega \in C_{\mathcal{F}}^{\circ}$, by Corollary 14, we deduce the following facts:

- ω is constant on each Q_u , hence $Q_u \subseteq F_i \setminus F_{i-1}$ for some $i = 1, \dots, |\mathcal{F}|$.
- if $Q_u, Q_v \subseteq F_i \setminus F_{i-1}$ for some $i = 1, \dots, |\mathcal{F}|$, then Q_u and Q_v are incomparable. Otherwise, if $i \leq_P j$ for some $i \in Q_u, j \in Q_v$, then ω^* is maximized along the ray generated by $e_i - e_j$, contrary to $\mathcal{F} \in F(C(Q))$.
- if $Q_u \subseteq F_i \setminus F_{i-1}, Q_v \subseteq F_j \setminus F_{j-1}$ and Q_u is smaller than Q_v , then $i <_Z j$. Otherwise, if $i >_Z j$, then $\omega_i > \omega_j$ for some $i \in Q_u, j \in Q_v$ such that $i <_P j$, hence ω^* does not reach the maximum.

We conclude that $\mathcal{F} \in \mathfrak{F}(P)$ and $P/\mathcal{F} = Q$, consequently $\text{rk}_P(\mathcal{F}) = n - m = \dim(C(Q))$.

⊆: Let $\mathcal{F} \triangleleft P$ be a flag of ideals of the poset P and $C : i_1, i_2, \dots, i_n$ be a circuit with all down-edges belonging to P/\mathcal{F} . The level function of elements of P according to the flag \mathcal{F} , defined by $l(i) = \min\{a \mid i \in F_a\}, i \in P$, is nondecreasing along the circuit C . It means that l is constant on C , i.e. $C \subseteq F_a \setminus F_{a-1}$, for some $1 \leq a \leq |\mathcal{F}|$. Particulary, all up-edges of the circuit C are in P/\mathcal{F} . The same is true for circuits with all up-edges in P/\mathcal{F} and we conclude that P/\mathcal{F} is a positive subposet of P . Corollary 14 gives $\mathcal{F} \in F(C(P/\mathcal{F}))$ and the proof is finished by the obvious identity $\text{rk}_P(\mathcal{F}) = \dim C(P/\mathcal{F})$. \square

By Theorem 21 and Proposition 17 we obtain the following expression for the f -polynomial.

Corollary 22. *Let $C(P)$ be the poset cone associated to a poset P on the ground set $[n]$. The f -polynomial of $C(P)$ is given by*

$$f(C(P), q) = (-1)^{n-1} \mathbf{ps}^1(\Psi_{-q}([P]))(-1).$$

Example 23. Let st_n be the poset on $[n]$ with covering relations $i \lessdot n$, for $1 \leq i \leq n-1$. Let $\mathcal{F} : \emptyset = F_0 \subset F_1 \subset \dots \subset F_k = [n]$ be a flag of ideals of the poset st_n . Then $n \in F_k$ and two different elements of $[n]$ are in the same connected component in st_n/\mathcal{F} if and only if both are in $F_k \setminus F_{k-1}$. Therefore

$$\text{rk}_{\text{st}_n}(\mathcal{F}) = |F_k \setminus F_{k-1}| - 1.$$

The number of flags corresponding to the faces of $C(\text{st}_n)$ of dimension i is equal to $\binom{n-1}{i}$, which implies

$$F_q(C(\text{st}_n)) = \sum_{i=0}^{n-1} \binom{n-1}{i} \left(M_{(1)}^{n-1-i} \right)_{i+1} q^i,$$

where for $F \in \mathcal{QSym}$, $F \mapsto (F)_i$ is the linear extension of the map given on monomial basis by $M_\alpha \mapsto M_{(\alpha, i)}$. By Proposition 22, the corresponding f -polynomial is equal to

$$f(C(\text{st}_n), q) = \sum_{i=0}^{n-1} \binom{n-1}{i} q^i = (1+q)^{n-1}.$$

Example 24. Let l_n be a linear poset on $[n]$ and $\mathcal{F} \in \mathfrak{F}(l_n)$. For $1 \leq i \leq |\mathcal{F}|$ all components of $F_i \setminus F_{i-1}$ are connected, so

$$c(F_i \setminus F_{i-1}) = 1 \text{ and } \text{rk}(\mathbb{1}_{F_{i+1} \setminus F_i}) = |F_{i+1} \setminus F_i| - 1.$$

It implies that $\text{rk}_{l_n}(\mathcal{F}) = n - |\mathcal{F}|$. We have

$$F_q(C(l_n)) = \sum_{i=0}^{n-1} \left(\sum_{\alpha: k(\alpha)=n-i} M_\alpha \right) q^i.$$

Since $|\{\alpha \models n : k(\alpha) = n - i\}| = \binom{n-1}{i}$, the corresponding f -polynomial is equal to

$$f(C(l_n), q) = \sum_{i=0}^{n-1} \binom{n-1}{i} q^i = (1+q)^{n-1}.$$

We obtain that $f(C(\text{st}_n), q) = f(C(l_n), q)$. Actually, it is a consequence of a more general fact.

Proposition 25. *Let \mathbf{P} be a poset on $[n]$ whose Hasse diagram is a tree. Then*

$$f(C(\mathbf{P}), q) = (1+q)^{n-1}.$$

Proof. The poset \mathbf{P} has $n - 1$ covering relations. Proposition 12 implies that the generating rays of $C(\mathbf{P})$ are $n - 1$ linearly independent vectors $e_i - e_j$ where $j < i$. Hence, the k -face of $C(\mathbf{P})$ is generated by k generating rays, so $f_k(C(\mathbf{P})) = \binom{n-1}{k}$. \square

Example 26. Let $\mathbf{K}_{m,n}$ be the poset on the set $[m+n]$ such that for all $i \in [m]$ and $j \in [m+n] \setminus [m]$ hold $i < j$. The Hasse diagram of $\mathbf{K}_{m,n}$ is the complete bipartite graph $K_{m,n}$. We have

$$\begin{aligned} F_q(C(\mathbf{K}_{m,n})) &= \left(M_{(1)}^m \right) \circ \left(M_{(1)}^n \right) + \\ &+ \sum_{k=1}^{m+n-1} q^k \sum_{t_1+t_2=k+1} \binom{m}{t_1} \binom{n}{t_2} \left(M_{(1)}^{m-t_1} \circ M_{(k+1)} \circ M_{(1)}^{n-t_2} \right), \end{aligned}$$

where $1 \leq t_1 \leq m$, $1 \leq t_2 \leq n$ and \circ is the *concatenation product* defined on monomial basis by $M_\alpha \circ M_\beta := M_{\alpha \cdot \beta}$. Note that this includes the case $\text{st}_n = \mathbf{K}_{n-1,1}$. The principal specialization evaluated at -1 gives

$$f(C(\mathbf{K}_{m,n}, q) = 1 + \sum_{k=1}^{m+n-1} q^k \sum_{t_1+t_2=k+1} \binom{m}{t_1} \binom{n}{t_2}.$$

4.4 Opposite poset

For a flag $\mathcal{F} : \emptyset = F_0 \subset F_1 \subset \cdots \subset F_k = [n]$, the *opposite flag* \mathcal{F}^{op} is defined by

$$\mathcal{F}^{op} : \emptyset \subset [n] \setminus F_{k-1} \subset [n] \setminus F_{k-2} \subset \cdots \subset [n] \setminus F_1 \subset [n].$$

The normal cone corresponding to the opposite flag is the opposite cone

$$C_{\mathcal{F}^{op}} = -C_{\mathcal{F}} \text{ and } \text{type}(\mathcal{F}^{op}) = \text{rev}(\text{type}(\mathcal{F})),$$

where $\text{rev}(\alpha_1, \alpha_2, \dots, \alpha_k) = (\alpha_k, \alpha_{k-1}, \dots, \alpha_1)$. The *opposite poset* \mathbf{P}^{op} to a poset \mathbf{P} is the poset on the same set such that $i <_{\mathbf{P}^{op}} j$ if and only if $j <_{\mathbf{P}} i$.

Proposition 27. *Let \mathbf{P} be a poset on $[n]$, the quasisymmetric enumerator function corresponding to the opposite poset \mathbf{P}^{op} is determined by*

$$F_q(C(\mathbf{P}^{op})) = \text{rev}(F_q(C(\mathbf{P}))),$$

where for $F \in \mathcal{QSym}$, $F \mapsto \text{rev}(F)$ is the linear extension of the map given on monomial basis by $M_\alpha \mapsto M_{\text{rev}(\alpha)}$.

Proof. Note that $\mathcal{F} \in \mathfrak{F}(\mathbf{P})$ if and only if $\mathcal{F}^{op} \in \mathfrak{F}(\mathbf{P}^{op})$. The statement follows from $\text{rk}_{\mathbf{P}}(\mathcal{F}) = \text{rk}_{\mathbf{P}^{op}}(\mathcal{F}^{op})$. \square

4.5 Action of the antipode on $F_q(C(\mathbf{P}))$

The geometric interpretation of the action of the antipode on \mathcal{QSym} is given by the following lemma.

Lemma 28 ([8], Lemma 4.5). *The antipode S on the monomial quasisymmetric function $M_{\mathcal{F}}$ associated to a flag \mathcal{F} acts by*

$$S(M_{\mathcal{F}}) = (-1)^{|\mathcal{F}|+1} \sum_{\mathcal{G} \preceq \mathcal{F}^{op}} M_{\mathcal{G}},$$

where $\mathcal{G} \preceq \mathcal{F}^{op}$ if and only if $\mathcal{F}^{op} \subseteq \mathcal{G}$ as faces of Pe^{n-1} .

The next theorem extends the similar statement proven for generalized permutohedra in [8] to the case of poset cones.

Theorem 29. *If \mathbf{P} is a poset on the set $[n]$, the antipode S acts on the quasisymmetric enumerator function $F_q(C(\mathbf{P}))$ by*

$$S(F_q(C(\mathbf{P}))) = (-1)^n \sum_{\mathcal{G} \in \mathfrak{F}(\mathbf{P})} f(C(\mathbf{P}/\mathcal{G}), -q) M_{\mathcal{G}^{op}}.$$

Proof. Since $\mathcal{F} \in \mathfrak{F}(\mathbf{P})$ and $\mathcal{G} \preceq \mathcal{F}$ implies $\mathcal{G} \in \mathfrak{F}(\mathbf{P})$ we have

$$\begin{aligned} S(F_q(C(\mathbf{P}))) &= \sum_{\mathcal{F} \in \mathfrak{F}(\mathbf{P})} q^{\mathrm{rk}_{\mathbf{P}}(\mathcal{F})} S(M_{\mathcal{F}}) = \sum_{\mathcal{F} \in \mathfrak{F}(\mathbf{P})} q^{\mathrm{rk}_{\mathbf{P}}(\mathcal{F})} (-1)^{|\mathcal{F}|+1} \sum_{\mathcal{G} \preceq \mathcal{F}^{op}} M_{\mathcal{G}} \\ &= \sum_{\mathcal{G} \in \mathfrak{F}(\mathbf{P}^{op})} M_{\mathcal{G}} \sum_{\substack{\mathcal{F} \in \mathfrak{F}(\mathbf{P}) \\ \mathcal{G} \preceq \mathcal{F}^{op}}} q^{\mathrm{rk}_{\mathbf{P}}(\mathcal{F})} (-1)^{|\mathcal{F}|+1}. \end{aligned}$$

By equivalences $\mathcal{G} \preceq \mathcal{F}^{op}$ if and only if $\mathcal{G}^{op} \preceq \mathcal{F}$ and $\mathcal{G} \in \mathfrak{F}(\mathbf{P}^{op})$ if and only if $\mathcal{G}^{op} \in \mathfrak{F}(\mathbf{P})$, the last equality becomes

$$\begin{aligned} S(F_q(C(\mathbf{P}))) &= (-1)^n \sum_{\mathcal{G}^{op} \in \mathfrak{F}(\mathbf{P})} M_{\mathcal{G}} \sum_{\substack{\mathcal{F} \in \mathfrak{F}(\mathbf{P}) \\ \mathcal{G}^{op} \preceq \mathcal{F}}} (-q)^{\mathrm{rk}_{\mathbf{P}}(\mathcal{F})} (-1)^{|\mathcal{F}|+1+n+\mathrm{rk}_{\mathbf{P}}(\mathcal{F})} \\ &= (-1)^n \sum_{\mathcal{G} \in \mathfrak{F}(\mathbf{P})} M_{\mathcal{G}^{op}} \sum_{\substack{\mathcal{F} \in \mathfrak{F}(\mathbf{P}) \\ \mathcal{G} \preceq \mathcal{F}}} (-q)^{\mathrm{rk}_{\mathbf{P}}(\mathcal{F})} (-1)^{|\mathcal{F}|+1+n+\mathrm{rk}_{\mathbf{P}}(\mathcal{F})}. \end{aligned}$$

By the proof of Theorem 21, we have that \mathbf{P}/\mathcal{G} is positive subposet of \mathbf{P} and \mathcal{G} is a normal flag to the face $C(\mathbf{P}/\mathcal{G})$. By the same argument, the faces of $C(\mathbf{P}/\mathcal{G})$ are of the form $C(\mathbf{P}/\mathcal{F})$ for flags of ideals $\mathcal{F} \in \mathfrak{F}(\mathbf{P})$ which satisfy $\mathcal{G} \preceq \mathcal{F}$. Therefore, according to the identities (1) and $\mathrm{rk}_{\mathbf{P}}(\mathcal{F}) = \dim C(\mathbf{P}/\mathcal{F})$, we obtain

$$f(C(\mathbf{P}/\mathcal{G}), q) = \sum_{\substack{\mathcal{F} \in \mathfrak{F}(\mathbf{P}) \\ \mathcal{G} \preceq \mathcal{F}}} (-1)^{n+1+|\mathcal{F}|+\mathrm{rk}_{\mathbf{P}}(\mathcal{F})} q^{\mathrm{rk}_{\mathbf{P}}(\mathcal{F})}.$$

□

5. THE FUNCTION $F(\mathbf{P})$

In this section we show that for a well labelled poset the enumerator function $F_q(C(\mathbf{P}))$ specializes at $q = 0$ to the enumerator of \mathbf{P} -partitions. For a poset \mathbf{P} on the set $[n]$ let

$$F(\mathbf{P}) = F_0(C(\mathbf{P})) = \sum_{\substack{\mathcal{F} \in \mathfrak{F}(\mathbf{P}) \\ \mathbf{P}/\mathcal{F} \text{ discrete}}} M_{\mathcal{F}}.$$

The sum is over flags of ideals of \mathbf{P} such that \mathbf{P}/\mathcal{F} is a *discrete poset* (poset with no covering relations).

Proposition 30. *The principal specialization \mathbf{ps}^1 of $F(\mathbf{P})$ at $m = -1$ results in*

$$\mathbf{ps}^1(F(\mathbf{P}))(-1) = (-1)^{n-1}.$$

Proof. The poset cone $C(\mathbf{P})$ has a unique vertex, so the required equation follows from Proposition 17. □

Let $P_1 * P_2$ be the series composition of posets P_1, P_2 which is a poset on the disjoint union of elements of P_1, P_2 with the order relation defined by $i \leq_{P_1 * P_2} j$ if and only if $i \leq_{P_1} j$ or $i \leq_{P_2} j$ or $i \in P_1, j \in P_2$.

Proposition 31. *If $P = P_1 * P_2 * \dots * P_n$ then*

$$F(P) = F(P_1) \circ F(P_2) \circ \dots \circ F(P_n),$$

where \circ is the concatenation product described in Example 26.

Proof. The proof follows from the fact that $F(P)$ is the sum over all flags of ideals \mathcal{F} such that P/\mathcal{F} is a discrete poset. Such a flag is the consecutive composition of flags of the same type corresponding to components P_1, \dots, P_n . \square

Example 32. A poset $K_{m,n}$ can be expressed as the series composition of discrete posets $d(m) * d(n)$ on $[m], [n]$. By Proposition 31, we obtain $F(K_{m,n}) = F(d(m)) \circ F(d(n)) = M_{(1)}^m \circ M_{(1)}^n$.

Proposition 33. *If $\text{Max}(P)$ is the set of maximal elements of a connected poset P on the set $[n]$ then*

$$F(P) = \sum_{\emptyset \neq A \subseteq \text{Max}(P)} (F(P|_{[n] \setminus A}))_{|A|}.$$

Particulary, if $\text{Max}(P) = \{v\}$ then $F(P) = (F(P|_{[n] \setminus \{v\}}))_1$.


Proof. For a flag of ideals $\mathcal{F} : \emptyset = F_0 \subset F_1 \subset \dots \subset F_m = [n]$ let $F_{\text{end}} = F_m \setminus F_{m-1}$. If P/\mathcal{F} is a discrete poset, we have that $P|_{F_{\text{end}}}$ is discrete too, so $F_{\text{end}} \subseteq \text{Max}(P)$. Therefore

$$F(P) = \sum_{\substack{\mathcal{F} \in \mathfrak{F}(P) \\ P/\mathcal{F} \text{ discrete}}} M_{\mathcal{F}} = \sum_{\emptyset \neq A \subseteq \text{Max}(P)} \sum_{\substack{\mathcal{F} \in \mathfrak{F}(P) \\ P/\mathcal{F} \text{ discrete} \\ F_{\text{end}} = A}} M_{\mathcal{F}} = \sum_{\emptyset \neq A \subseteq \text{Max}(P)} (F(P|_{[n] \setminus A}))_{|A|}.$$

\square

Remark 34. The direct calculation shows that $F(P)$ distinguishes all non-isomorphic posets on $[n]$, for $n \in \{1, 2, 3, 4, 5, 6\}$.

The weighted enumerator function invariant F_q contains more information about posets than its specialization at $q = 0$. The following example of posets with the same quasisymmetric invariant $F(P)$ is borrowed from [9, Example 4.9].

Figure 7: Posets with the same quasisymmetric invariant $F(P)$

Example 35. Let P_1 and P_2 be posets on Figure 7. The direct calculation gives

$$\begin{aligned}
F(P_1) = F(P_2) = & M_{(2,3,2)} + 2M_{(1,1,3,2)} + 2M_{(2,3,1,1)} \\
& + M_{(1,3,2,1)} + M_{(1,2,3,1)} + M_{(2,1,3,1)} + M_{(1,3,1,2)} \\
& + 3M_{(1,2,2,2)} + 3M_{(2,1,2,2)} + 3M_{(2,2,1,2)} + 3M_{(2,2,2,1)} \\
& + 2M_{(1,2,2,1,1)} + 6M_{(2,1,1,1,2)} + 6M_{(1,2,2,1,1)} \\
& + 7M_{(2,1,1,2,1)} + 7M_{(1,2,1,1,2)} + 8M_{(1,1,2,1,2)} + 8M_{(2,1,2,1,1)} \\
& + 8M_{(1,2,1,2,1)} + 8M_{(1,1,2,2,1)} + 9M_{(1,1,1,2,2)} + 9M_{(2,2,1,1,1)} \\
& + 20M_{(1,1,1,1,1,2)} + 23M_{(1,1,1,1,2,1)} + 24M_{(1,1,1,2,1,1)} \\
& + 24M_{(1,1,2,1,1,1)} + 23M_{(1,2,1,1,1,1)} + 20M_{(2,1,1,1,1,1)} \\
& + 4M_{(1,1,3,1,1)} + 3M_{(1,3,1,1,1)} + 3M_{(1,1,1,3,1)} + 66M_{(1,1,1,1,1,1,1)}.
\end{aligned}$$

Consider flags $\emptyset \subset \{i\} \subset [6] \setminus \{j\} \subset [6]$, for $i \in \{1, 2\}$ and $j \in \{5, 6\}$ which are the only flags of ideals of the type $(1, 5, 1)$ of posets P_1 and P_2 . The coefficients by $M_{(1,5,1)}$ in $F_q(C(P_i))$, $i = 1, 2$ are

$$\zeta_{(1,5,1)}(P_1) = 2q^4 + q^3 + q^2 \quad \text{and} \quad \zeta_{(1,5,1)}(P_2) = q^4 + 3q^3,$$

which shows that $F_q(C(P_1)) \neq F_q(C(P_2))$.

The following theorem gives a geometric interpretation of the enumerator $F_P(\mathbf{x})$ of P -partitions for a well labelled poset P . Recall from Definition 2 that

$$F_P(\mathbf{x}) = \sum_{f \in \mathcal{A}(P)} x_{f(1)} x_{f(2)} \cdots x_{f(n)},$$

where $f \in \mathcal{A}(P)$ if and only if $i <_P j$ implies $f(i) < f(j)$, for all $i, j \in P$.

Proposition 36. *For a well labelled poset P it holds that*

$$\mathcal{A}(P) = \{\omega \in C_{\mathcal{F}}^{\circ} : \mathcal{F} \in \mathfrak{F}(P) \text{ and } P/\mathcal{F} \text{ is a discrete poset}\}.$$

Proof. We have $\omega \in C_{\mathcal{F}}^{\circ}$ for some $\mathcal{F} \in \mathfrak{F}(P)$ such that P/\mathcal{F} is a discrete poset if and only if ω^* is maximized uniquely at the vertex of $C(P)$. By Proposition 13 and definition of P -partitions of well labeled posets, both sets are characterized by the same condition. \square

Theorem 37. For a well labelled poset P it holds that $F(P) = F_P(\mathbf{x})$.

Proof. By (3) and Proposition 36

$$F(P) = \sum_{\substack{\mathcal{F} \in \mathfrak{F}(P) \\ P/\mathcal{F} \text{ discrete}}} M_{\mathcal{F}} = \sum_{\substack{\omega \in \mathbb{Z}_+^n \cap C_{\mathcal{F}}^{\circ} \\ \mathcal{F} \in \mathfrak{F}(P) \\ P/\mathcal{F} \text{ discrete}}} x_{\omega_1} x_{\omega_2} \cdots x_{\omega_n} = \sum_{\omega \in \mathcal{A}(P)} x_{\omega_1} x_{\omega_2} \cdots x_{\omega_n},$$

which is exactly the expression for $F_P(\mathbf{x})$. \square

Acknowledgements. The authors are thankful to anonymous referees for careful reading and valuable comments that improved the manuscript.

REFERENCES

1. M. AGUIAR, F. ARDILA: *Hopf monoids and generalized permutahedra*, [arXiv:1709.07504](#)
2. M. AGUIAR, N. BERGERON AND F. SOTTILE: *Combinatorial Hopf algebras and generalized Dehn-Sommerville relations*, *Compositio Math.* **142** (2006), 1–30.
3. L. BILLERA, N. JIA, V. REINER: *A quasisymmetric function for matroids*, *European J. Comb.* **30** (2009) 1727–1757.
4. L. BILLERA, A. SARANGARAJAN: *The combinatorics of permutation polytopes*, *Formal power series and algebraic combinatorics*, *Series formelles et combinatoires algebriques* (1994)
5. D. GRINBERG, V. REINER: *Hopf Algebras in Combinatorics*, [arXiv:1409.8356](#).
6. V. GRUJIĆ: *Quasisymmetric functions for nestohedra*, *SIAM J. Discrete Math.* **31**(4) (2017) 2570–2585.
7. V. GRUJIĆ: *Counting faces of graphical zonotopes*, *Ars Math. Contemp.* **13** (2017) 227–234.
8. V. GRUJIĆ, M. PEŠOVIĆ, T. STOJADINOVIĆ: *Weighted quasisymmetric enumerator for generalized permutohedra*, *J. Algebr. Comb.* (2019) [doi:10.1007/s10801-019-00874-x](#)
9. P. R. W. MCNAMARA AND R. E. WARD: *Equality of P-partition generating functions*, *Ann. Comb.* **18**(3) (2014) 489–514.
10. A. POSTNIKOV: *Permutohedra, associahedra, and beyond*, *Int. Math. Res. Not.* **6** (2009), 1026–1106.
11. A. POSTNIKOV, V. REINER AND L. WILLIAMS: *Faces of generalized permutohedra*, *Documenta Math.* **13** (2008), 207–273.

Marko Pešović,
University of Belgrade, Faculty of Civil Engineering,
Bulevar kralja Aleksandra 73,
11000 Belgrade, Serbia
E-mail: mpesovic@grf.bg.ac.rs

(Received 25. 05. 2020.)

(Revised 08. 04. 2021.)

Tanja Stojadinović,

University of Belgrade, Faculty of Mathematics,

Studentski trg 16,

11000 Belgrade, Serbia

E-mail: *tanjas@matf.bg.ac.rs*