

WHAT IS THE BEST FRACTIONAL DERIVATIVE TO FIT DATA?

Ricardo Almeida

The aim of this work is to show, based on concrete data observation, that the choice of the fractional derivative when modelling a problem is relevant for the accuracy of a method. Using the least squares fitting technique, we determine the order of the fractional differential equation that better describes the experimental data, for different types of fractional derivatives.

1. INTRODUCTION

Fractional differential equations (FDE) are an extension of ODE's, where the order of the derivative may take any real positive value [6, 9]. For this reason, very often we can model an experimental dynamic more efficiently by considering the problem formulated as a FDE. For example, they have found applications in viscoelastic [17], mechanics [3, 7], economy [10], signal processing [15], biology [1], logistic population models [14], etc. A question that always arises when doing so is what fractional derivative is to be considered, since there are various definitions in the literature [11]. Usually, the answer to this question depends on the problem and on the observed data. One of the most common definition, at least for real world applications, is the Caputo fractional derivative. Given a function $f \in C^n[a, b]$ and a positive real α , let $n \in \mathbb{N}$ be such that $\alpha \in (n - 1, n)$. The Caputo fractional derivative of f of order α is given by the integral

$${}^C D_{a+}^{\alpha} f(x) := \frac{1}{\Gamma(n - \alpha)} \int_a^x (x - t)^{n - \alpha - 1} f^{(n)}(t) dt.$$

2010 Mathematics Subject Classification. 26A33, 34A08, 90C30.
Keywords and Phrases. Fractional calculus, modelling, optimization.

Here, Γ denotes the Gamma function:

$$\Gamma(x) := \int_0^\infty t^{x-1} e^{-t} dt, \quad x > 0.$$

This special function is an extension of the factorial function, in the sense that

$$\Gamma(x + 1) = x\Gamma(x), \quad x > 0 \quad \text{and} \quad \Gamma(k) = (k - 1)!, \quad k \in \mathbb{N}.$$

One way to overcome this vast number of definitions for fractional derivatives is to consider a more general definition for fractional operators, and then determine which ones fit better with the data [11]. Let $\alpha > 0$, $f \in L^1[a, b]$ and $\psi \in C^1[a, b]$ be an increasing function such that $\psi'(x) \neq 0$, for all $x \in [a, b]$. The fractional integral of f with respect to ψ is defined as

$$I_{a+}^{\alpha, \psi} f(x) := \frac{1}{\Gamma(\alpha)} \int_a^x \psi'(t) (\psi(x) - \psi(t))^{\alpha-1} f(t) dt.$$

Fractional derivatives are defined in the following way. If $f, \psi \in C^n[a, b]$ (with ψ increasing and $\psi'(x) \neq 0$, for all x), the fractional derivative of f with respect to ψ is given by [2]

$${}^C D_{a+}^{\alpha, \psi} f(x) := \frac{1}{\Gamma(n - \alpha)} \int_a^x \psi'(t) (\psi(x) - \psi(t))^{n-\alpha-1} \left(\frac{1}{\psi'(t)} \frac{d}{dt} \right)^n f(t) dt,$$

where $\alpha \in (n - 1, n)$ and $n \in \mathbb{N}$. We recall the Mittag-Leffler function E_α : given $\alpha > 0$ and $x \in \mathbb{R}$, this special function is given by the series

$$E_\alpha(x) := \sum_{k=0}^\infty \frac{x^k}{\Gamma(\alpha k + 1)}.$$

It can be seen as a generalization of the exponential function, since when $\alpha = 1$, we have

$$E_\alpha(x) = \exp(x).$$

For example, we have the following:

$$\begin{aligned} {}^C D_{a+}^{\alpha, \psi} (\psi(x) - \psi(a))^{\beta-1} &= \frac{\Gamma(\beta)}{\Gamma(\beta - \alpha)} (\psi(x) - \psi(a))^{\beta-\alpha-1}, \quad \beta > n, \quad \text{and} \\ {}^C D_{a+}^{\alpha, \psi} E_\alpha(\lambda(\psi(x) - \psi(a))^\alpha) &= \lambda E_\alpha(\lambda(\psi(x) - \psi(a))^\alpha), \quad \lambda \in \mathbb{R}. \end{aligned}$$

Fractional integral and fractional derivative are the inverse operation of each other, in the sense that

$$I_{a+}^{\alpha, \psi} {}^C D_{a+}^{\alpha, \psi} f(x) = f(x) - \sum_{k=0}^{n-1} \frac{\left(\frac{1}{\psi'(x)} \frac{d}{dx} \right)^k f(a)}{k!} (\psi(x) - \psi(a))^k$$

and

$$I_{a+}^{\alpha, \psi C} D_{a+}^{\alpha, \psi} f(x) = f(x) - f(a).$$

Based on experimental data, our goal is to test several FDE's and see which ones better describe the problems. All the numerical computations are done in MAPLE, using the *Statistics* package and the Maple tool *NonlinearFit*. This tool fits a nonlinear model to a given data, by minimizing the least-squares error. Suppose that the n points $(\tilde{x}_i, \tilde{y}_i)$ are the original data and the model has the form $F(A, x)$, where the parameter A is a m dimensional vector. The objective is to find the values of the parameters for the model such that the squared residuals

$$\text{Error} = \sum_{i=1}^n (\tilde{y}_i - F(A, \tilde{x}_i))^2$$

is a minimum. In each section, we will fit three kinds of models to the data points. First, we consider the classical model, which corresponds to the case $\alpha = 1$. Secondly, we consider the method to be fit when the process is described by the Caputo fractional derivative, that is, when we take $\psi_1(x) := x$. In the third model, we consider three different kernels $\psi_2(x) := (x + 1)^b$, $\psi_3(x) := \ln(x + 1)$ and $\psi_4(x) := \sin(x/b)$, with $b > 0$, and choose the one that better fits with the original data. We test the efficiency of the model, by comparing the error of the classical model, Error_C , with the error given by the fractional model, Error_F :

$$\text{Efficiency} = \frac{\text{Error}_C - \text{Error}_F}{\text{Error}_C} \times 100.$$

We will study four problems: the Gross Domestic Product (Section), Newton's law of cooling (Section), the Bombay Plague Epidemic (Section), and the World population growth (Section). The problems considered are described by a linear differential equation because when we replace after the ordinary derivative by a fractional derivative, we know the exact solution to the fractional differential equation.

2. GROSS DOMESTIC PRODUCT

The Gross Domestic Product (GDP) is one of the most important of all economic statistics, used to measure the performance of a country's economy. It represents the total of finished goods and services produced over a specific time period. We consider in this section the GDP of two countries: the USA and the UK. We will describe them by a linear and by an exponential model:

$$S'(t) = k \quad \text{and} \quad S'(t) = kS(t).$$

If we consider the model described by the FDE's,

$${}^C D_{0+}^{\alpha, \psi} S(t) = k \quad \text{or by} \quad {}^C D_{0+}^{\alpha, \psi} S(t) = kS(t),$$

with $\alpha \in (0, 1)$, the solutions are given by

$$S(t) = S_0 + I_{0+}^{\alpha, \psi} k \quad \text{and} \quad S(t) = S_0 E_{\alpha}(k(\psi(t) - \psi(0))^{\alpha}),$$

respectively. To get a better accuracy for the model, we will allow the fractional order α to be arbitrary, and not only on the open interval $(0, 1)$. For the data, we use the ones available from the World Bank [16], which consist in GDP per capita between the years 1966 until 2011, counted every 5 years. The variable t , in years, starts at $t = 0$, corresponding to the year 1966, and $S(T)$ is the GDP per capita, in US\$. In Tables 1 and 2 we present the results obtained when considering the USA case, with linear and exponential growth, and in Tables 3 and 4, the UK case.

Model	k	α	Error	Efficiency
Classical	938.653334	1	1.163673×10^8	
Fractional ψ_1	251.518736	1.439934	9.895013×10^6	91.496740%
Fractional ψ_4 ($b = 50$)	1.050515	1.664986	7.346859×10^6	93.686491%

Table 1: GDP in the USA: linear case.

Model	k	α	Error	Efficiency
Classical	0.058556	1	2.302460×10^8	
Fractional ψ_1	0.267899	0.400291	4.129788×10^7	82.063587%
Fractional ψ_3	1.501965	4.880700	8.241729×10^6	96.420468%

Table 2: GDP in the USA: exponential case.

Model	k	α	Error	Efficiency
Classical	785.672510	1	2.315927×10^8	
Fractional ψ_1	105.319149	1.683547	7.114080×10^7	69.687165%
Fractional ψ_3	9.274772×10^4	5.961180	6.947902×10^7	70.408906%

Table 3: GDP in the UK: linear case.

Model	k	α	Error	Efficiency
Classical	0.071713	1	2.562247×10^8	
Fractional ψ_1	0.359075	0.33255	9.980382×10^7	61.048327%
Fractional ψ_3	4.412682	5.873892	6.951255×10^7	72.870476%

Table 4: GDP in the UK: exponential case.

Either way, the best model to describe the problem is when we consider linear growth. This allows us to predict the GDP in future years more accurately. For example, in the year 2014, the GDP in the USA is about 54629\$ and in the UK 46332\$. Using the classical model and the values of Tables 1 and 3, the predictions are 49202\$ for the USA, and 39672\$ for the UK. So, the errors are

$$54629 - 49202 = 5427 \quad \text{and} \quad 46332 - 39672 = 6660,$$

respectively. When considering the fractional models, the predictions are 54305\$ for the USA and 47616\$ for the UK. The errors are now

$$54629 - 54305 = 324 \quad \text{and} \quad 47616 - 46332 = 1284,$$

respectively. In Figure 1 we show the optimal curves, for the USA and UK cases, considering the linear and exponential growths.




Figure 1: Gross Domestic Product in the USA and UK.

3. NEWTON'S LAW OF COOLING

Newton's law of cooling states that the rate of change of the temperature of a body is directly proportional to the difference between its own temperature and the ambient temperature, provided the difference is small. This statement leads to

the classic equation of exponential decline over time, which can be applied to many phenomena. Newton was able to show that temperature change follows the ODE

$$T'(t) = k(T(t) - T_a),$$

where $T(t)$ is the temperature of the body at a given time t , T_a is the ambient temperature, and k is a cooling constant, specific to the object. The solution of this EDO is the exponential function

$$T(t) = T_a + (T_0 - T_a) \exp(kt),$$

where T_0 is the starting temperature of the object. If we consider the problem modeled by a FDE of order $\alpha > 0$, then we get that the solution of the equation

$${}^C D_{0+}^{\alpha, \psi} T(t) = k(T(t) - T_a)$$

is the function

$$T(t) = T_a + (T_0 - T_a) E_{\alpha}(k(\psi(t) - \psi(0))^{\alpha}).$$

To test the accuracy of the model an experiment was carried out. With a beaker filled with 100ml of water, originally at $100^{\circ}C$, and with a thermometers in the beaker, the temperature of the water was measured every minute. The ambient temperature for this investigation was $23^{\circ}C$, and the results obtained can be viewed in [5]. In Table 5 we present the numerical results for this problem, and in Figure 2 we compare the plots for the classical model with the two fractional models.

Model	k	α	Error	Efficiency
Classical	-0.0665349	1	940.640679	
Fractional ψ_1	-0.130333	0.798038	39.844721	95.764087%
Fractional ψ_3	-0.249423	1.428019	26.691909	97.162369%

Table 5: Newton’s law of cooling.

4. BOMBAY PLAGUE EPIDEMIC, 1905-6

Between 1896 and 1914, India was struck by a series of major epidemics: malaria, cholera, Spanish influenza, and the bubonic plague. In Bombay City alone, the number of deaths were up to 183984 on that period. Kermack and McKendrick proposed a model [8] based on the number of deaths from October 1905 until September 1906 [4, Table IX], counted every week. If $R(t)$ denotes the number of deaths in the week t , for $t \in \{0, \dots, 51\}$, with $t = 0$ corresponding to 1st of October 1905, the authors proposed the model obtained from the ODE

$$R'(t) = A / \cosh^2(Bt + C).$$

The solution is given by the function

$$R(t) = R_0 + \int_0^x \frac{A}{\cosh^2(B\tau + C)} d\tau.$$




Figure 2: Newton's law of cooling.

When we consider the fractional model,

$${}^C D_{0+}^{\alpha, \psi} R(t) = A / \cosh^2(Bt + C),$$

with $\alpha \in (0, 1)$, applying the fractional integral to both sides of the equation, we deduce that

$$\begin{aligned} R(t) &= R_0 + I_{0+}^{\alpha, \psi} \frac{A}{\cosh^2(Bt + C)} \\ &= R_0 + \frac{1}{\Gamma(\alpha)} \int_0^x \psi'(\tau) (\psi(t) - \psi(\tau))^{\alpha-1} \frac{A}{\cosh^2(B\tau + C)} d\tau. \end{aligned}$$

The results obtained are presented in Table 6. In Figure 3 we show the plots for the total number of deaths, and for the number of deaths per week. To obtain a better model we will consider $\alpha > 0$. As we can see, when we model the problem using the Caputo fractional derivative the gain in efficiency is very small. However, considering another kernel we obtain a gain of approximately 5%.

Model	A	B	C	α	Error	Efficiency
Classical	736.714057	0.183346	-5.070156	1	6.309364×10^5	
Fractional ψ_1	733.259286	0.184230	-5.088524	1.002834	6.302366×10^5	0.110908%
Fractional ψ_3	2.217880×10^4	0.194156	-5.478100	1.017019	5.983701×10^5	5.161572%

Table 6: Bombay Plague Epidemic.

5. WORLD POPULATION GROWTH

Historically, the most important law to characterize a population growth is described by the differential equation

$$N'(t) = \lambda N(t),$$




Figure 3: Bombay Plague Epidemic.

where λ is the population growth rate. The solution to this problem, with N_0 the initial number of individuals, is the exponential function

$$N(t) = N_0 \exp(\lambda t).$$

When we consider the problem modeled by a FDE, that is, by the equation

$${}^C D_{0+}^{\alpha, \psi} N(t) = \lambda N(t),$$

the solution is given by

$$N(t) = N_0 E_{\alpha}(\lambda(\psi(t) - \psi(0))^{\alpha}).$$

We study the optimal curve that better fits with the data in four continents: Africa, America, Asia and Europe. For data source, we use the one available from the UN [12]. It consists of the number of individuals in the several continents, from 1950 until 2010, counted every 5 years. The initial value $t = 0$ corresponds to the year 1950, with $N(t)$ measured in millions. In Figure 4 and in Tables 7, 8, 9 and 10, we show the results.

Model	λ	α	Error	Efficiency
Classical	0.025116	1	327.024434	
Fractional ψ_1	0.022421	1.03793	236.643618	27.637328%
Fractional ψ_2 ($b = 0.7$)	0.027381	1.699578	40.953972	87.476785%

Table 7: World population: Africa.

Model	λ	α	Error	Efficiency
Classical	0.017806	1	6660.898290	
Fractional ψ_1	0.042814	0.723857	685.697720	89.705627%
Fractional ψ_4 ($b = 60$)	1.114633	0.915357	10.987758	99.835041%

Table 8: World population: America.




Figure 4: World population.

Model	λ	α	Error	Efficiency
Classical	0.019087	1	1.378920×10^9	
Fractional ψ_1	0.040394	0.761743	49095.37114	64.395774%
Fractional ψ_4 ($b = 50$)	1.326809	1.124085	920.911794	99.332150%

Table 9: World population: Asia.

Model	λ	α	Error	Efficiency
Classical	0.005879	1	6246.195248	
Fractional ψ_1	0.035060	0.485031	776.467956	87.568945%
Fractional ψ_4 ($b = 40$)	0.263725	0.711280	155.33140	97.513184%

Table 10: World population: Europe.

For comparison, in Table 11, we present the classical and the best fractional model, for the year 2015, with the estimated population in each of the continents

analysed previously [13].

Continent	Population	Classical	Error	Fractional	Error
Africa	1166	1171	5	1161	5
America	991	1080	89	990	1
Asia	4384	4826	442	4342	42
Europe	743	805	62	743	0

Table 11: World population in 2015.

Acknowledgments. Work supported by Portuguese funds through the CIDMA - Center for Research and Development in Mathematics and Applications, and the Portuguese Foundation for Science and Technology (FCT-Fundação para a Ciência e a Tecnologia), within project UID/MAT/04106/2013.

REFERENCES

1. R. ALMEIDA, N.R.O. BASTOS, M.T.T. MONTEIRO: *Modelling some real phenomena by fractional differential equations*. Math. Meth. Appl. Sci., **39** (16) (2016), 4846–4855.
2. R. ALMEIDA: *A Caputo fractional derivative of a function with respect to another function*. Commun. Nonlinear Sci. Numer. Simul., **44** (2017), 460–481.
3. A. CARPINTERI, P. CORNETTI, A. SAPORA: *Nonlocal elasticity: an approach based on fractional calculus*. Meccanica, **49** (11) (2014), 2551–2569.
4. M. GREENWOOD: *XXII. The epidemiological observations made by the commission in Bombay city. Appendix II. Report on certain plague statistics*. J. Hygiene 7 (1907) 794–798.
5. E. GIESEKING: *Newton's law of cooling. An experimental investigation (2014)*. <http://jwilson.coe.uga.edu/EMAT6680Fa2014/Giesecking/Exploration%2012/Newton's%20Law%20of%20Cooling.htm>. Accessed 14 March 2016.
6. TADEUSZ JANKOWSKI: *Positive solutions to advanced fractional differential equations with nonlocal boundary conditions*. Appl. Anal. Discrete Math., **9** (2015), 209–220.
7. J. T. KATSIKADELIS: *Generalized fractional derivatives and their applications to mechanical systems*. Arch. Appl. Mech., **85** (9) (2014), 1307–1320.
8. W.O. KERMAK, A.G. MCKENDRICK: *Contribution to the Mathematical Theory of Epidemics*. Proceedings of the Royal Society of London. Series A, **115** (1927), 700–721.
9. BARBARA ŁUPIŃSKA, TATIANA ODZIJEWICZ, EWA SCHMEIDEL: *On the solutions to a generalized fractional cauchy problem*. Appl. Anal. Discrete Math., **10** (2016), 332–344.
10. J.A.T. MACHADO, M.E. MATA: *A fractional perspective to the bond graph modelling of world economies*. Nonlinear Dyn., **80** (4) (2014), 1839–1852.
11. S. G. SAMKO, A. A. KILBAS, O. I. MARICHEV: *Fractional integrals and derivatives, translated from the 1987 Russian original*. Gordon and Breach, Yverdon, 1993.

12. UNITED NATIONS: *Department of Economic and Social Affairs, Population Division (2013)*. World Population Prospects. http://www.geohive.com/earth/his_proj_continent.aspx. Accessed 8 March 2016.
13. UNITED NATIONS: *Department of Economic and Social Affairs, Population Division*. Population Projection for 2015. http://www.nationsonline.org/oneworld/world_population.htm. Accessed 8 March 2016.
14. Ş YÜZBAŞI: *A collocation method for numerical solutions of fractional-order logistic population model*. Int. J. Biomath., **9 (2)** (2016), 1650031.
15. J. WANG, Y. YE, Y. GAO, S. QIAN, X. GAO: *Fractional Compound Integral with Application to ECG Signal Denoising*. Circuits Systems Signal Process., **34 (6)** (2014), 1915–1930.
16. WORLD BANK: *World Development Indicators*. <http://databank.worldbank.org/data/reports.aspx?source=2&country=&series=NY.GDP.PCAP.CD&period=>. Accessed 16 March 2016.
17. J.M.P. ZERPA, A. CANELAS, B. SENSALÉ, D.B. SANTANA, R.L. ARMENTANO: *Modeling the arterial wall mechanics using a novel high-order viscoelastic fractional element*. Appl. Math. Model., **39 (16)** (2015), 4767–4780.

Ricardo Almeida

Center for Research and Development
in Mathematics and Applications (CIDMA)
Department of Mathematics,
University of Aveiro,
3810–193 Aveiro, Portugal
E-mail: ricardo.almeida@ua.pt

(Received 26.07.2016.)

(Revised 04.03.2017.)