

**NEW FAMILIES OF SPECIAL NUMBERS FOR  
COMPUTING NEGATIVE ORDER EULER NUMBERS  
AND RELATED NUMBERS AND POLYNOMIALS**

*Yilmaz Simsek*

The main purpose of this paper is to construct new families of special numbers with their generating functions. These numbers are related to many well-known numbers, which are Bernoulli numbers, Fibonacci numbers, Lucas numbers, Stirling numbers of the second kind and central factorial numbers. Our other inspiration of this paper is related to the Golombek's problem [15] "Aufgabe 1088. El. Math., 49 (1994), 126–127". Our first numbers are not only related to the Golombek's problem, but also computation of the negative order Euler numbers. We compute a few values of the numbers which are given by tables. We give some applications in probability and statistics. That is, special values of mathematical expectation of the binomial distribution and the Bernstein polynomials give us the value of our numbers. Taking derivative of our generating functions, we give partial differential equations and also functional equations. By using these equations, we derive recurrence relations and some formulas of our numbers. Moreover, we come up with a conjecture with two open questions related to our new numbers. We give two algorithms for computation of our numbers. We also give some combinatorial applications, further remarks on our new numbers and their generating functions.

---

2010 Mathematics Subject Classification. 11B83, 12D10, 11B68, 11S40, 11S80, 26C05, 26C10, 30B40, 30C15.

Keywords and Phrases. Fibonacci numbers; Bernoulli numbers; Euler numbers; Central factorial numbers; Array polynomials; Stirling numbers; Generating functions; Functional equations; Binomial coefficients; Combinatorial sum.

## 1. INTRODUCTION

In this section, we consider the following question:

What could be more basic tools to compute the negative order of the first and the second kind Euler numbers? One of motivations of this paper is associated with this question and its answer. Another motivation of this paper is related to the work of Golombek [15], which is entitled **Aufgabe 1088**.

Here, let  $\mathbb{C}$ ,  $\mathbb{R}$ ,  $\mathbb{Q}$ ,  $\mathbb{Z}$ ,  $\mathbb{N}$  be the sets of complex numbers, real numbers, rational numbers, integers, and positive integers, respectively and let  $\mathbb{N}_0 = \{0, 1, 2, 3, \dots\} = \mathbb{N} \cup \{0\}$ .

Golombek gave the following novel combinatorial sum:

$$(1) \quad \sum_{j=0}^k \binom{k}{j} j^n = \frac{d^n}{dt^n} (e^t + 1)^k \Big|_{t=0},$$

where  $n \in \mathbb{N}$ . Golombek [15] also mentioned that this sum is related to the following sequence

$$n2^{n-1}, n(n+1)2^{n-2}, \dots$$

We introduce new families of special numbers, which are not only used in counting techniques and problems, but also computing negative order of the first and the second kind Euler numbers and other combinatorial sums. Here, our technique is related to the generating functions and their functional equations. In the historical development of mathematics, we can observe that the generating functions play a very important role in pure and applied mathematics. These function are powerful tools in solving counting problems and investigating properties of the special numbers and polynomials. In addition, the generating functions are also used in computer programming, in physics, and in other areas. Briefly, in Physics, generating functions, which arise in Hamiltonian mechanics, are quite different from generating functions in mathematics. The generating functions are functions whose partial derivatives generate the differential equations that determine a system's dynamics. These functions are also related to the partition function of statistical mechanics (cf. [11], [21], [37]). In mathematics, a generating function can be expanded as formal power series in one indeterminate whose coefficients encode information about a sequence of numbers and that is indexed by the natural numbers (cf. [11], [12], [14], [13], [21], [37], [29], [40]). As far as we know, the generating function is firstly discovered by Abraham de Moivre (26 May 1667 -27 November 1754, French mathematician) (cf. [21]). In order to solve the general linear recurrence problem, Moivre constructed the concept of the generating functions in 1730. In work of Doubilet et al. [14], we also see that Laplace (23 March 1749-5 March 1827, French mathematician, physicist and statistician) discovered the remarkable correspondence between set theoretic operations and operations on formal power series. Their method gives us great success to solve a variety of combinatorial problems. They developed new kinds of algebras of generating functions better suited to combinatorial and probabilistic problems. Their method depends on group algebra

(or semigroup algebra) (see, for details, [14]). It is well-known that there are many different ways or approaches to generate a sequence of numbers and polynomials from the series or the generating functions. The purpose of this paper is to construct the generating functions for new families of numbers involving Golombek's identity in (1), Stirling numbers, central factorial numbers, Euler numbers of negative order, rook numbers and combinatorial sums. Our method and approach provides a way of constructing new special families of numbers and combinatorial sums. We show how several of these numbers and these combinatorial sums relate to each other. We pose a conjecture with two open questions associated with our new numbers and their generating functions.

We organize our paper as follows:

In Section 2, we briefly review some special numbers and polynomials, which are Bernoulli numbers, Euler numbers, Stirling numbers, central factorial numbers and array polynomials.

In Section 3, we give a generating function. By using this function, we define a family of new numbers  $y_1(n, k; \lambda)$ . We investigate many properties including recurrence relations of these numbers by using their generating functions. We compute a few values of the numbers  $y_1(n, k; \lambda)$ , which are given by tables. We give some remarks and comments related to the Golombek's identity and the numbers  $y_1(n, k; 1)$ . Finally, we give a conjecture with two open questions.

In Section 4, we give a generating function for a new family of the other numbers  $y_2(n, k; \lambda)$ . By using this function, we investigate many properties with a recurrence relation of these numbers. We compute a few values of the numbers  $y_2(n, k; \lambda)$ , which are given by tables. We give relations between these numbers, Fibonacci numbers, Lucas numbers, and  $\lambda$ -Stirling numbers of the second kind. We also give some combinatorial sums.

In Section 5, we define  $\lambda$ -central factorial numbers  $C(n, k; \lambda)$ . By using their generating function, we derive some identities and relations including these numbers and the others.

In Section 6, we give some applications related to the special values of mathematical expectation for the binomial distribution, the Bernstein polynomials and the Bernoulli polynomials.

In Section 7, by using the numbers  $y_1(n, k; \lambda)$ , we compute the Euler numbers of negative order. In addition, we compute a few values of these numbers, which are given by tables.

In Section 8, we give two algorithms for our computations.

In Section 9, we give combinatorial applications, including a rook numbers and polynomials. We also give combinatorial interpretation for the numbers  $y_1(n, k; 1)$ . Finally in the last section, we give further remarks with conclusion.

The principal value  $\ln z$  is the logarithm whose imaginary part lies in the interval  $(-\pi, \pi]$ . Moreover we also use the following notational conventions:

$$0^n = \begin{cases} 1, & (n = 0) \\ 0, & (n \in \mathbb{N}) \end{cases}$$

and

$$\binom{\lambda}{0} = 1 \text{ and } \binom{\lambda}{v} = \frac{\lambda(\lambda-1)\cdots(\lambda-v+1)}{v!} = \frac{(\lambda)_v}{v!} \quad (n \in \mathbb{N}, \lambda \in \mathbb{C})$$

(cf. [4], [12], [42]). For combinatorial example, we will use the notations of Bona [5], that is the set  $\{1, 2, \dots, n\}$  is an  $n$ -element set, that is,  $n$  distinct objects. Therefore, Bona introduced the shorter notation  $[n]$  for this set. The number  $n(n-1)(n-2)\cdots(n-k+1)$  of all  $k$ -element lists from  $[n]$  without repetition occurs so often in combinatorics that there is a symbol for it, namely

$$(n)_k = n(n-1)(n-2)\cdots(n-k+1)$$

(cf. [5, pp. 11-13.]).

## 2. Background

In this section, we give a brief introduction about Bernoulli numbers, Euler numbers, the  $(\lambda)$ -Stirling numbers and array polynomials, which will be used in subsequent sections.

In [2]-[45], we see that there are many known properties and relations involving various kind of the special numbers and polynomials such as Bernoulli polynomials and numbers, Euler polynomials and numbers, Stirling numbers and also rook polynomials and numbers by making use of some standard techniques based upon generating functions and other known techniques.

Bernoulli polynomials are defined by means of the following generating function (cf. [13]-[45]):

$$\frac{t}{e^t - 1} e^{tx} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}$$

( $|t| < 2\pi$ ). One can observe that

$$B_n = B_n(0),$$

which denotes Bernoulli numbers (cf. [13]-[45]; see also the references cited in each of these earlier works).

The sum of powers of integers is related to the Bernoulli numbers and polynomials:

$$(2) \quad \sum_{k=0}^n k^r = \frac{1}{r+1} (B_{r+1}(n+1) - B_{r+1}),$$

(cf. [13], [40], [42]).

The first kind Apostol-Euler polynomials of order  $k$ , with  $k \geq 0$ ,  $E_n^{(k)}(x; \lambda)$  are defined by means of the following generating function:

$$(3) \quad F_{P_1}(t, x; k, \lambda) = \left( \frac{2}{\lambda e^t + 1} \right)^k e^{tx} = \sum_{n=0}^{\infty} E_n^{(k)}(x; \lambda) \frac{t^n}{n!}$$

( $|t| < \pi$  when  $\lambda = 1$  and  $|t| < |\ln(-\lambda)|$  when  $\lambda \neq 1$ ),  $\lambda \in \mathbb{C}$ ,  $k \in \mathbb{N}$  with, of course,

$$E_n^{(k)}(\lambda) = E_n^{(k)}(0; \lambda),$$

which denote the first kind Apostol-Euler numbers of order  $k$  (cf. [20], [13], [28], [26], [30], [40], [45]). Substituting  $k = \lambda = 1$  into (3), we have the first kind Euler numbers  $E_n = E_n^{(1)}(1)$ , which are defined by means of the following generating function:

$$\frac{2}{e^t + 1} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!},$$

( $|t| < \pi$ ) (cf. [13]-[45]; see also the references cited in each of these earlier works).

The second kind Euler numbers  $E_n^*$  are defined by means of the following generating function:

$$\frac{2}{e^t + e^{-t}} = \sum_{n=0}^{\infty} E_n^* \frac{t^n}{n!}$$

( $|t| < \frac{\pi}{2}$ ) (cf. [8], [13], [24], [26], [30], [42], [45]; see also the references cited in each of these earlier works).

Stirling numbers of the second kind are used in pure and applied mathematics. These numbers occur in combinatorics and in the theory of partitions. The Stirling numbers of the second kind, denoted by  $S_2(n, v)$ , the number of ways to partition a set of  $n$  objects into  $k$  groups ([5], [7], [12], [37], [42]).

Let  $v \in \mathbb{N}_0$  and  $\lambda \in \mathbb{C}$ . The  $\lambda$ -Stirling numbers of the second kind  $S_2(n, v; \lambda)$  are generalized of the Stirling number of the second kind. These numbers  $S_2(n, v; \lambda)$  are defined by means of the following generating function:

$$(4) \quad F_S(t, v; \lambda) = \frac{(\lambda e^t - 1)^v}{v!} = \sum_{n=0}^{\infty} S_2(n, v; \lambda) \frac{t^n}{n!}.$$

For further information about these numbers, the reader may be referred to [25] and ([34], [33], [39]; see also the references cited in each of these earlier works).

Observe that

$$S_2(n, v) = S_2(n, v; 1),$$

which are computing by the following formulas:

$$x^n = \sum_{v=0}^n \binom{x}{v} v! S_2(n, v)$$

or

$$S_2(n, v) = \frac{1}{v!} \sum_{j=0}^v \binom{v}{j} (-1)^j (v-j)^n$$

(cf. [13]-[45]; see also the references cited in each of these earlier works). A recurrence relation for these numbers is given by

$$S_2(n, k) = S_2(n-1, k-1) + kS_2(n-1, k),$$

with

$$S_2(n, 0) = 0 \ (n \in \mathbb{N}); \ S_2(n, n) = 1 \ (n \in \mathbb{N}); \ S_2(n, 1) = 1 \ (n \in \mathbb{N})$$

and  $S_2(n, k) = 0$  ( $n < k$  or  $k < 0$ ) (cf. [13]-[45]; see also the references cited in each of these earlier works).

Let  $v \in \mathbb{N}_0$  and  $\lambda \in \mathbb{C}$ . In [34], we defined the  $\lambda$ -array polynomials  $S_v^n(x; \lambda)$  by means of the following generating function:

$$(5) \quad F_A(t, x, v; \lambda) = \frac{(\lambda e^t - 1)^v}{v!} e^{tx} = \sum_{n=0}^{\infty} S_v^n(x; \lambda) \frac{t^n}{n!}$$

(cf. [4], [34]).

The array polynomials  $S_v^n(x)$  are defined by means of the following generating function:

$$(6) \quad F_A(t, x, v) = \frac{(e^t - 1)^v}{v!} e^{tx} = \sum_{n=0}^{\infty} S_v^n(x) \frac{t^n}{n!},$$

(cf. [4], [10], [34]; see also the references cited in each of these earlier works). By using the above generating function, we have

$$S_v^n(x) = \frac{1}{v!} \sum_{j=0}^v (-1)^{v-j} \binom{v}{j} (x+j)^n$$

with

$$S_0^0(x) = S_n^n(x) = 1, \ S_0^n(x) = x^n$$

and for  $v > n$ ,

$$S_v^n(x) = 0$$

(cf. [10], [34], [35]; see also the references cited in each of these earlier works).

Recently, central factorial numbers  $T(n, k)$  have been studied by many authors. These numbers are used in theory of numbers, combinatorics and probability. Central factorial numbers  $T(n, k)$  (of the second kind) are defined by means of the following generating function:

$$(7) \quad F_T(t, k) = \frac{1}{(2k)!} (e^t + e^{-t} - 2)^k = \sum_{n=0}^{\infty} T(n, k) \frac{t^{2n}}{(2n)!}$$

(cf. [5], [7], [12], [41], [35], [37]; see also the references cited in each of these earlier works).

These numbers have the following relations:

$$x^n = \sum_{k=0}^n T(n, k)x(x-1)(x-2^2)(x-3^2)\cdots(x-(k-1)^2).$$

Combining the above equation with (7), we also have

$$T(n, k) = T(n-1, k-1) + k^2T(n-1, k),$$

where  $n \geq 1$ ,  $k \geq 1$ ,  $(n, k) \neq (1, 1)$ . For  $n, k \in \mathbb{N}$ ,  $T(0, k) = T(n, 0) = 0$  and  $T(n, 1) = 1$  (cf. [5], [7], [12], [41], [35], [37]).

### 3. A family of new numbers $y_1(n, k; \lambda)$

In this section, we give generating function for the numbers  $y_1(n, k; \lambda)$ . We give some functional equations and differential equations of this generating function. By using these equations, we derive various new identities and combinatorial relations involving these numbers. Some of our observations on these numbers can be briefly expressed as follows: the numbers  $y_1(n, k; \lambda)$  are related to the  $\lambda$ -Stirling numbers of the second kind, the central factorial numbers, the Euler numbers of negative orders and the Golombek's identity.

Let  $k \in \mathbb{N}_0$  and  $\lambda \in \mathbb{C}$ . We define these numbers,  $y_1(n, k; \lambda)$  by the following generating function:

$$(8) \quad F_{y_1}(t, k; \lambda) = \frac{1}{k!} (\lambda e^t + 1)^k = \sum_{n=0}^{\infty} y_1(n, k; \lambda) \frac{t^n}{n!}.$$

The function  $F_{y_1}(t, k; \lambda)$  is an analytic function.

By using (8), we get

$$\sum_{n=0}^{\infty} y_1(n, k; \lambda) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left( \frac{1}{k!} \sum_{j=0}^k \binom{k}{j} j^n \lambda^j \right) \frac{t^n}{n!}.$$

Comparing the coefficients of  $t^n$  on both sides of the above equation, we arrive at the the following theorem:

**Theorem 1.** *Let  $n \in \mathbb{N}_0$ . The following identity holds:*

$$(9) \quad y_1(n, k; \lambda) = \frac{1}{k!} \sum_{j=0}^k \binom{k}{j} j^n \lambda^j.$$

We assume that  $\lambda \neq 0$ . For  $k = 0, 1, 2, 3, 4$  and  $n = 0, 1, 2, 3, 4, 5$ , we use (9) to compute a few values of the numbers  $y_1(n, k; \lambda)$  as follows:

$n \setminus k$	0	1	2	3	4
0	1	$\lambda + 1$	$\frac{1}{2}\lambda^2 + \lambda + \frac{1}{2}$	$\frac{1}{6}\lambda^3 + \frac{1}{2}\lambda^2 + \frac{1}{2}\lambda + \frac{1}{6}$	$\frac{1}{24}\lambda^4 + \frac{1}{6}\lambda^3 + \frac{1}{4}\lambda^2 + \frac{1}{6}\lambda + \frac{1}{24}$
1	0	$\lambda$	$\lambda^2 + \lambda$	$\frac{1}{2}\lambda^3 + \lambda^2 + \frac{1}{2}\lambda$	$\frac{1}{6}\lambda^4 + \frac{1}{2}\lambda^3 + \frac{1}{2}\lambda^2 + \frac{1}{6}\lambda$
2	0	$\lambda$	$2\lambda^2 + \lambda$	$\frac{3}{2}\lambda^3 + 2\lambda^2 + \frac{1}{2}\lambda$	$\frac{5}{3}\lambda^4 + \frac{3}{2}\lambda^3 + \lambda^2 + \frac{1}{6}\lambda$
3	0	$\lambda$	$4\lambda^2 + \lambda$	$\frac{9}{2}\lambda^3 + 4\lambda^2 + \frac{1}{2}\lambda$	$\frac{8}{3}\lambda^4 + \frac{9}{2}\lambda^3 + 2\lambda^2 + \frac{1}{6}\lambda$
4	0	$\lambda$	$8\lambda^2 + \lambda$	$\frac{27}{2}\lambda^3 + 8\lambda^2 + \frac{1}{2}\lambda$	$\frac{32}{3}\lambda^4 + \frac{27}{2}\lambda^3 + 4\lambda^2 + \frac{1}{6}\lambda$
5	0	$\lambda$	$16\lambda^2 + \lambda$	$\frac{81}{2}\lambda^3 + 16\lambda^2 + \frac{1}{2}\lambda$	$\frac{128}{3}\lambda^4 + \frac{81}{2}\lambda^3 + 8\lambda^2 + \frac{1}{6}\lambda$

**Table 1:** Some numerical values of the numbers  $y_1(n, k; \lambda)$ .

For  $k = 0, 1, 2, \dots, 9$  and  $n = 0, 1, 2, \dots, 9$ , we also use (9) to compute a few values of the numbers  $y_1(n, k; 1)$  as follows:

$n \setminus k$	0	1	2	3	4	5	6	7	8	9
0	1	2	2	$\frac{4}{3}$	$\frac{2}{3}$	$\frac{4}{15}$	$\frac{4}{45}$	$\frac{8}{315}$	$\frac{2}{315}$	$\frac{4}{2835}$
1	0	1	2	$\frac{4}{3}$	$\frac{2}{3}$	$\frac{4}{15}$	$\frac{4}{45}$	$\frac{8}{315}$	$\frac{2}{315}$	$\frac{4}{2835}$
2	0	1	3	4	$\frac{10}{3}$	2	$\frac{14}{15}$	$\frac{16}{45}$	$\frac{4}{35}$	$\frac{2}{63}$
3	0	1	5	9	$\frac{28}{3}$	$\frac{20}{3}$	$\frac{18}{5}$	$\frac{14}{9}$	$\frac{176}{315}$	$\frac{6}{35}$
4	0	1	9	22	$\frac{85}{3}$	24	$\frac{224}{15}$	$\frac{328}{45}$	$\frac{102}{35}$	$\frac{62}{63}$
5	0	1	17	57	$\frac{274}{3}$	$\frac{275}{3}$	$\frac{328}{5}$	$\frac{1624}{45}$	$\frac{5048}{315}$	$\frac{208}{35}$
6	0	1	33	154	$\frac{925}{3}$	367	$\frac{4529}{15}$	$\frac{8416}{45}$	$\frac{3224}{35}$	$\frac{2360}{63}$
7	0	1	65	429	$\frac{3238}{3}$	$\frac{4580}{3}$	$\frac{7223}{5}$	$\frac{9065}{9}$	$\frac{173216}{315}$	$\frac{8576}{35}$
8	0	1	129	1222	$\frac{11665}{3}$	6554	$\frac{107114}{15}$	$\frac{252268}{45}$	$\frac{118717}{35}$	$\frac{104288}{63}$
9	0	1	257	3537	$\frac{42994}{3}$	$\frac{86645}{3}$	$\frac{181458}{5}$	$\frac{1444534}{45}$	$\frac{6781748}{315}$	$\frac{402723}{35}$

**Table 2:** Some numerical values of the numbers  $y_1(n, k; 1)$ .

Some special values of  $y_1(n, k; \lambda)$  are given as follows:

$$y_1(0, k; \lambda) = \frac{1}{k!}(\lambda + 1)^k,$$

$$y_1(n, 0; \lambda) = 0, \quad (n \in \mathbb{N})$$

and

$$y_1(n, 1; \lambda) = \lambda, \quad (n \in \mathbb{N}).$$

By using (8), we derive the following functional equation

$$\lambda^k e^{kt} = \sum_{l=0}^k (-1)^{k-l} \binom{k}{l} l! F_{y_1}(t, l; \lambda).$$



Combining (8) with the above equation, we get

$$\lambda^k \sum_{n=0}^{\infty} \frac{(kt)^n}{n!} = \sum_{n=0}^{\infty} \left( \sum_{l=0}^k (-1)^{k-l} \binom{k}{l} l! y_1(n, l; \lambda) \right) \frac{t^n}{n!}.$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we obtain the following theorem:

**Theorem 2.**

$$k^n \lambda^k = \sum_{l=0}^k (-1)^{k-l} \binom{k}{l} l! y_1(n, l; \lambda).$$

We give a relationship between the numbers  $y_1(n, k; \lambda)$  and the  $\lambda$ -Stirling numbers of the second kind by the following theorem:

**Theorem 3.**

$$S_2(n, k; \lambda^2) = \frac{k!}{2^n} \sum_{l=0}^n \binom{n}{l} S_2(l, k; \lambda) y_1(n-l, k; \lambda).$$

*Proof.* By using (4) and (8), we derive the following functional equation:

$$F_S(2t, k; \lambda^2) = k! F_S(t, k; \lambda) F_{y_1}(t, k; \lambda).$$

From the above equation, we have

$$\sum_{n=0}^{\infty} 2^n S_2(n, k; \lambda^2) \frac{t^n}{n!} = \sum_{n=0}^{\infty} S_2(n, k; \lambda) \frac{t^n}{n!} \sum_{n=0}^{\infty} y_1(n, k; \lambda) \frac{t^n}{n!}.$$

Therefore

$$\sum_{n=0}^{\infty} 2^n S_2(n, k; \lambda^2) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left( k! \sum_{l=0}^n \binom{n}{l} S_2(l, k; \lambda) y_1(n-l, k; \lambda) \right) \frac{t^n}{n!}.$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we get the desired result.  $\square$

A relationship between the numbers  $y_1(n, k; \lambda)$ ,  $S_2(n, k; \lambda^3)$  and the array polynomials  $S_k^n(x; \lambda)$  is given by the following theorem:

**Theorem 4.**

$$S_2(n, k; \lambda^3) = \sum_{l=0}^n \sum_{j=0}^k \binom{n}{l} \binom{k}{j} \frac{\lambda^{2k-2j} j!}{3^n} y_1(l, j; \lambda) S_k^{n-l}(2k-2j; \lambda).$$

*Proof.* Combining (4), (5) and (8), we get

$$F_S(3t, k; \lambda^3) = \sum_{j=0}^k \frac{k!}{(k-j)!} \lambda^{2k-2j} F_A(t, 2k-2j, k; \lambda) F_{y_1}(t, j; \lambda).$$

By using the above functional equation, we obtain

$$\sum_{n=0}^{\infty} 3^n S_2(n, k; \lambda^3) \frac{t^n}{n!} = \sum_{j=0}^k \frac{k!}{(k-j)!} \lambda^{2k-2j} \sum_{n=0}^{\infty} S_k^n(n, 2k-2j; \lambda) \frac{t^n}{n!} \sum_{n=0}^{\infty} y_1(n, j; \lambda) \frac{t^n}{n!}.$$

Therefore

$$\sum_{n=0}^{\infty} 3^n S_2(n, k; \lambda^3) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \sum_{l=0}^n \sum_{j=0}^k \binom{n}{l} \binom{k}{j} j! \lambda^{2k-2j} y_1(l, j; \lambda) S_k^{n-l}(2k-2j; \lambda) \frac{t^n}{n!}.$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we arrive at the desired result.  $\square$

There are many combinatorial applications for (9). That is, by substituting  $\lambda = 1$  into (9), we set

$$(10) \quad B(n, k) = k! y_1(n, k; 1).$$

In [15], Golombek gave the following formula for (9):

$$B(n, k) = \frac{d^n}{dt^n} (e^t + 1)^k \Big|_{t=0}.$$

**Remark 1.** If we substitute  $\lambda = -1$  into (9), then we get the Stirling numbers of the second kind (cf. [5]-[44]):

$$S_2(n, k) = (-1)^k y_1(n, k; -1) = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n.$$

**Remark 2.** We claim that the numbers  $B(n, k)$  are related to the following numbers:

$$a_k 2^k$$

where the sequence  $a_k$  is a positive integer depend on  $k$ . Consequently, in the work of Spivey Identity 8-Identity 10 Spevy, we see

$$B(0, k) = 2^k,$$

$$B(1, k) = k 2^{k-1},$$

$$B(2, k) = k(k+1) 2^{k-2},$$

see also [5, p. 56, Exercise 21] and [11, p. 117].

**Remark 3.** In [38, Identity 12.], Spivey also proved the following novel identity by the falling factorial method:

$$(11) \quad B(m, n) = \sum_{j=0}^m \binom{n}{j} j! 2^{n-j} S_2(m, j).$$

The numbers  $B(0, k)$  are given by means of the following well-known generating function: Let  $|x| < \frac{1}{2}$ , we have

$$\sum_{k=0}^{\infty} B(0, k) x^k = \frac{1}{1-2x}.$$

The numbers  $B(1, k)$  are given by means of the following well-known generating function: Let  $|x| < \frac{1}{2}$ , we have

$$\sum_{k=1}^{\infty} B(1, k) x^k = \frac{x}{(1-2x)^2}.$$

**Remark 4.** In work of Boyadzhiev [6, p.4, Eq-(7)], we have

$$\sum_{j=0}^k \binom{k}{j} j^n x^j = \sum_{j=0}^n \binom{k}{j} j! S_2(n, j) x^j (1+x)^{k-j}.$$

Substituting  $x = 1$  into the above equation, we arrive at (11).

**Theorem 5.** Let  $d \in \mathbb{N}$  and  $m_0, m_1, m_2, \dots, m_d \in \mathbb{Q}$ . Let  $m_0 \neq 0$ . Thus we have

$$(12) \quad \sum_{v=0}^{d-1} m_v B(d-v, k) = 2^{k-d} \binom{k}{d}.$$

*Proof.* It is well-known that

$$(1+x)^k = \sum_{j=0}^k \binom{k}{j} x^j.$$

Taking the  $d^{\text{th}}$  derivative, with respect to  $x$ , we obtain

$$(13) \quad \binom{k}{d} (1+x)^{k-d} = \sum_{j=0}^k \binom{k}{j} \binom{j}{d} x^{j-d}.$$

Substituting  $x = 1$  into the above equation, we get

$$(14) \quad 2^{k-d} \binom{k}{d} = \sum_{j=0}^k \binom{k}{j} \binom{j}{d}.$$

In [36], we have

$$\binom{j}{d} = m_0 j^d + m_1 j^{d-1} + \cdots + m_{d-1} j,$$

where  $m_0, m_1, \dots, m_{d-1} \in \mathbb{Q}$ . Therefore

$$2^{k-d} \binom{k}{d} = \sum_{j=0}^k \binom{k}{j} (m_0 j^d + m_1 j^{d-1} + \cdots + m_{d-1} j).$$

Thus we get

$$2^{k-d} \binom{k}{d} = \sum_{v=0}^{d-1} m_v \sum_{j=0}^k \binom{k}{j} j^{d-v}.$$

Combining (10) with the above equation, we have

$$2^{k-d} \binom{k}{d} = \sum_{v=0}^{d-1} m_v B(d-v, k).$$

This completes the proof.  $\square$

There are many combinatorial arguments of (13). That is, if we substitute  $d = 3$  and  $4$  into (13), then we compute  $B(3, k)$  and  $B(4, k)$ , respectively, as follows:

$$B(3, k) = k^2(k+3)2^{k-3}$$

and

$$B(4, k) = k(k^3 + 6k^2 + 3k - 2)2^{k-4}.$$

By using (12), we derive the following result:

$$B(d, k) = \frac{2^{k-d}}{m_0} \binom{k}{d} - \sum_{v=1}^{d-1} \frac{m_v}{m_0} B(d-v, k).$$

Therefore, we *conjecture* that

$$B(d, k) = (k^d + x_1 k^{d-1} + x_2 k^{d-2} + \cdots + x_{d-2} k^2 + x_{d-1} k) 2^{k-d},$$

where  $x_1, x_2, \dots, x_{d-1}$  and  $d$  are positive integers. Consequently, we arrive at the following open questions:

- 1-How can we compute the coefficients  $x_1, x_2, \dots, x_{d-1}$ ?
- 2-We assume that for  $|x| < r$

$$\sum_{k=1}^{\infty} B(d, k) x^k = f_d(x).$$

Is it possible to find  $f_d(x)$  function?

### 3.1. Recurrence relation and some identities for the numbers $y_1(n, k; \lambda)$

Here, by applying derivative operator to the generating functions (8), we give a recurrence relation and other formulas for the numbers  $y_1(n, k; \lambda)$ .

**Theorem 6.** *Let  $k \in \mathbb{N}$ . The following identity holds:*

$$y_1(n+1, k; \lambda) = ky_1(n, k; \lambda) - y_1(n, k-1; \lambda).$$

*Proof.* Taking derivative of (8), with respect to  $t$ , we obtain the following partial differential equation:

$$\frac{\partial}{\partial t} F_{y_1}(t, k; \lambda) = kF_{y_1}(t, k; \lambda) - F_{y_1}(t, k-1; \lambda).$$

Combining (8) with the above equation, we get

$$\sum_{n=1}^{\infty} y_1(n, k; \lambda) \frac{t^{n-1}}{(n-1)!} = k \sum_{n=0}^{\infty} y_1(n, k; \lambda) \frac{t^n}{n!} - \sum_{n=0}^{\infty} y_1(n, k-1; \lambda) \frac{t^n}{n!}.$$

After some elementary calculations, comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we obtain the desired result.  $\square$

**Theorem 7.** *Let  $k \in \mathbb{N}$ . The following identity holds:*

$$\frac{\partial}{\partial \lambda} y_1(n, k; \lambda) = \sum_{j=0}^n \binom{n}{j} y_1(j, k-1; \lambda).$$

*Proof.* Taking derivative of (8), with respect to  $\lambda$ , we obtain the following partial differential equation:

$$(15) \quad \frac{\partial}{\partial \lambda} F_{y_1}(t, k; \lambda) = e^t F_{y_1}(t, k-1; \lambda).$$

Combining (8) with the above equation, we get

$$\sum_{n=0}^{\infty} \frac{\partial}{\partial \lambda} y_1(n, k; \lambda) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \sum_{j=0}^n \binom{n}{j} y_1(j, k-1; \lambda) \frac{t^n}{n!}.$$

After some elementary calculations, comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we obtain the desired result.  $\square$

**Theorem 8.** *Let  $k \in \mathbb{N}$ . The following identity holds:*

$$\lambda \frac{\partial}{\partial \lambda} y_1(n, k; \lambda) = ky_1(n, k; \lambda) - y_1(n, k-1; \lambda).$$

*Proof.* By using (15), we obtain the following partial differential equation:

$$\lambda \frac{\partial}{\partial \lambda} F_{y_1}(t, k; \lambda) = k F_{y_1}(t, k; \lambda) - F_{y_1}(t, k-1; \lambda).$$

Combining (8) with the above equation, we get

$$\sum_{n=0}^{\infty} \lambda \frac{\partial}{\partial \lambda} y_1(n, k; \lambda) \frac{t^n}{n!} = \sum_{n=0}^{\infty} k y_1(n, k; \lambda) \frac{t^n}{n!} - \sum_{n=0}^{\infty} y_1(n, k-1; \lambda) \frac{t^n}{n!}.$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we get the desired result.  $\square$

#### 4. A family of new numbers $y_2(n, k; \lambda)$

In this section, we define a family of new numbers  $y_2(n, k; \lambda)$  by means of the following generating function:

$$(16) \quad F_{y_2}(t, k; \lambda) = \frac{1}{(2k)!} (\lambda e^t + \lambda^{-1} e^{-t} + 2)^k = \sum_{n=0}^{\infty} y_2(n, k; \lambda) \frac{t^n}{n!},$$

where  $k \in \mathbb{N}_0$  and  $\lambda \in \mathbb{C}$ .

By using (16) with their functional equation, we derive various identities and relations including our new numbers, the Fibonacci numbers, the Lucas numbers, the Stirling numbers and the central factorial numbers.

We get the following explicit formula for the numbers  $y_2(n, k; \lambda)$ :

**Theorem 9.** *Let  $n, k \in \mathbb{N}$ . The following identity holds:*

$$(17) \quad y_2(n, k; \lambda) = \frac{1}{(2k)!} \sum_{j=0}^k \binom{k}{j} 2^{k-j} \sum_{l=0}^j \binom{j}{l} (2l-j)^n \lambda^{2l-j}.$$

*Proof.* By (16), we have

$$\sum_{n=0}^{\infty} y_2(n, k; \lambda) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left( \frac{1}{(2k)!} \sum_{j=0}^k \binom{k}{j} 2^{k-j} \sum_{l=0}^j \binom{j}{l} (2l-j)^n \lambda^{2l-j} \right) \frac{t^n}{n!}.$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we obtain the desired result.  $\square$

For  $k = 0, 1, 2, 3$  and  $n = 0, 1, 2, 3, 4, 5$ , we use (17) to compute a few values of the numbers  $y_2(n, k; \lambda)$  as follows:

$n \setminus k$	0	1	2	3
0	1	$\frac{1}{2\lambda} + \frac{\lambda}{2} + 1$	$\frac{\lambda^2+4\lambda}{24} + \frac{4\lambda+1}{24\lambda^2} + \frac{1}{4}$	$\frac{\lambda^3+6\lambda^2}{720} + \frac{\lambda}{48} + \frac{1}{48\lambda} + \frac{6\lambda+1}{720\lambda^3} + \frac{1}{36}$
1	0	$\frac{\lambda}{2} - \frac{1}{2\lambda}$	$\frac{\lambda^2+2\lambda}{12} - \frac{2\lambda+1}{6\lambda^2}$	$\frac{\lambda^3+4\lambda^2}{240} + \frac{\lambda}{48} - \frac{1}{48\lambda} - \frac{4\lambda+1}{240\lambda^3}$
2	0	$\frac{\lambda}{2} + \frac{1}{2\lambda}$	$\frac{\lambda^2+\lambda}{6} + \frac{\lambda+1}{6\lambda^2}$	$\frac{\lambda^3}{80} + \frac{\lambda^2}{30} + \frac{\lambda}{48} + \frac{1}{48\lambda} + \frac{1}{30\lambda^2} + \frac{1}{80\lambda^3}$
3	0	$\frac{\lambda}{2} - \frac{1}{2\lambda}$	$\frac{2\lambda^2+\lambda}{6} - \frac{\lambda+2}{6\lambda^2}$	$\frac{3\lambda^3}{80} + \frac{\lambda^2}{15} + \frac{\lambda}{48} - \frac{1}{48\lambda} - \frac{1}{15\lambda^2} - \frac{3}{80\lambda^3}$
4	0	$\frac{\lambda}{2} + \frac{1}{2\lambda}$	$\frac{2\lambda^2+\lambda}{3} + \frac{\lambda+4}{6\lambda^2}$	$\frac{9\lambda^3}{80} + \frac{2\lambda^2}{15} + \frac{\lambda}{48} + \frac{1}{48\lambda} + \frac{2}{15\lambda^2} + \frac{9}{80\lambda^3}$
5	0	$\frac{\lambda}{2} - \frac{1}{2\lambda}$	$\frac{8\lambda^2+\lambda}{6} - \frac{\lambda+8}{6\lambda^2}$	$\frac{27\lambda^3}{80} + \frac{4\lambda^2}{15} + \frac{\lambda}{48} - \frac{1}{48\lambda} - \frac{4}{15\lambda^2} - \frac{27}{80\lambda^3}$

**Table 3:** Some numerical values of the numbers  $y_2(n, k; \lambda)$ .

By using (8) and (16), we get the following functional equation:

$$F_{y_2}(t, k; \lambda) = \frac{k!}{(2k)!} \sum_{j=0}^k F_{y_1}(t, j; \lambda) F_{y_1}(-t, k-j; \lambda^{-1}).$$

By combining (8) and (16) with the above equation, we obtain

$$\sum_{n=0}^{\infty} y_2(n, k; \lambda) \frac{t^n}{n!} = \frac{k!}{(2k)!} \sum_{j=0}^k \left( \sum_{n=0}^{\infty} y_1(n, j; \lambda) \frac{t^n}{n!} \sum_{n=0}^{\infty} (-1)^n y_1(n, k-j; \lambda^{-1}) \frac{t^n}{n!} \right).$$

Therefore

$$\sum_{n=0}^{\infty} y_2(n, k; \lambda) \frac{t^n}{n!} = \frac{k!}{(2k)!} \sum_{n=0}^{\infty} \sum_{j=0}^k \sum_{l=0}^n (-1)^{n-l} \binom{n}{l} y_1(l, j; \lambda) y_1(n-l, k-j; \lambda^{-1}) \frac{t^n}{n!}.$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, the numbers  $y_2(n, k; \lambda)$  is given in terms of the numbers  $y_1(n, k; \lambda)$  by the following theorem:

**Theorem 10.** *The following identity holds:*

$$(18) \quad y_2(n, k; \lambda) = \frac{k!}{(2k)!} \sum_{j=0}^k \sum_{l=0}^n (-1)^{n-l} \binom{n}{l} y_1(l, j; \lambda) y_1(n-l, k-j; \lambda^{-1}).$$

**Theorem 11.** *The following identity holds:*

$$y_1(n, 2k; \lambda) = \lambda^k \sum_{j=0}^n \binom{n}{j} k^{n-j} y_2(j, k; \lambda).$$

*Proof.* By using (8) and (16), we get the following functional equation:

$$\lambda^k e^{kt} F_{y_2}(t, k; \lambda) = F_{y_1}(t, 2k; \lambda).$$

From the above functional equation, we obtain

$$\sum_{n=0}^{\infty} y_1(n, 2k; \lambda) \frac{t^n}{n!} = \lambda^k \sum_{n=0}^{\infty} \frac{(kt)^n}{n!} \sum_{n=0}^{\infty} y_2(n, k; \lambda) \frac{t^n}{n!}.$$

Therefore

$$\sum_{n=0}^{\infty} y_1(n, 2k; \lambda) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left( \lambda^k \sum_{j=0}^n \binom{n}{j} k^{n-j} y_2(j, k; \lambda) \right) \frac{t^n}{n!}.$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we obtain the desired result.  $\square$

By substituting  $\lambda = 1$  into (16), we have

$$F_{y_2}(t, k) = \frac{1}{(2k)!} (e^t + e^{-t} + 2)^k.$$

The function  $F_{y_2}(t, k)$  is an even function. Consequently, we get the following result:

$$y_2(2n + 1, k; 1) = 0.$$

Thus, we get

$$(19) \quad F_{y_2}(t, k) = F_{y_2}(t, k; 1) = \sum_{n=0}^{\infty} y_2(n, k) \frac{t^{2n}}{(2n)!}.$$

By using (19), we give the following explicit formula for the numbers  $(y_2(n, k) = y_2(n, k; 1))$ :

**Corollary 1.** *The following identity holds:*

$$(20) \quad y_2(n, k) = \frac{1}{(2k)!} \sum_{j=0}^k \binom{k}{j} 2^{k-j} \sum_{l=0}^j \binom{j}{l} (2l - j)^n.$$

From the equation (20), we see that

$$y_2(0, 0) = 1.$$

For  $k = 0, 1, 2, \dots, 9$  and  $n \in \mathbb{N}$ , we use (20) to compute a few values of the numbers  $y_2(n, k)$  as follows:

$$y_2(n, 0) = 0,$$

$$y_2(n, 1) = \frac{(-1)^n + 1}{2},$$



$$y_2(n, 2) = \frac{(-1)^n + 1}{6} + \frac{2^{n-3} - (-2)^{n-3}}{3},$$

$$y_2(n, 3) = \frac{(-1)^n + 1}{48} + \frac{2^{n-3} - (-2)^{n-3}}{15} + \frac{3^{n-2} + (-3)^{n-2}}{80},$$

$$y_2(n, 4) = \frac{(-1)^n + 1}{720} + \frac{2^{n-4} + (-2)^{n-4}}{105} + \frac{2^{n-5} - (-2)^{n-5}}{315} + \frac{3^{n-2} + (-3)^{n-2}}{560} + \frac{4^{n-3} - (-4)^{n-3}}{630},$$

$$y_2(n, 5) = \frac{(-1)^n + 1}{17280} + \frac{2^{n-4} + (-2)^{n-4} + 2^{n-5} - (-2)^{n-5}}{2835} + \frac{3^{n-2} + (-3)^{n-2}}{8960} + \frac{2(4^{n-4} + (-4)^{n-4})}{2835} + \frac{5^{n-2} + (-5)^{n-2}}{145152},$$

$$y_2(n, 6) = \frac{(-1)^n + 1}{604800} + \frac{2^{n-5} - (-2)^{n-5} + 2^{n-10} + (-2)^{n-10}}{31185} + \frac{3^{n-4} + (-3)^{n-4}}{98560} + \frac{3^{n-5} - (-3)^{n-5}}{12320} + \frac{4^{n-4} + (-4)^{n-4}}{31185} + \frac{2(4^{n-5} - (-4)^{n-5})}{155925} + \frac{5^{n-2} + (-5)^{n-2}}{1596672} + \frac{6^{n-5} - (-6)^{n-5}}{61600},$$

$$y_2(n, 7) = \frac{(-1)^n + 1}{29030400} + \frac{2^{n-3} - (-2)^{n-3}}{6081075} + \frac{2^{n-10} + (-2)^{n-10}}{405405} + \frac{3^{n-3} - (-3)^{n-3}}{7321600} + \frac{3^{n-5} - (-3)^{n-5}}{640640} + \frac{4^{n-4} + (-4)^{n-4}}{1216215} + \frac{2(4^{n-5} - (-4)^{n-5})}{2027025} + \frac{5^{n-2} + (-5)^{n-2}}{38320128} + \frac{6^{n-5} - (-6)^{n-5}}{800800} + \frac{7^{n-2} + (-7)^{n-2}}{1779148800},$$

$$y_2(n, 8) = \frac{(-1)^n + 1}{1828915200} + \frac{7^{n-2} + (-7)^{n-2}}{26687232000} + \frac{5^{n-2} + (-5)^{n-2}}{1494484992} + \frac{3^{n-4} + (-3)^{n-4}}{64064000} + \frac{4^{n-5} - (-4)^{n-5}}{18243225} + \frac{8^{n-5} - (-8)^{n-5}}{638512875} + \frac{4^{n-5} - (-4)^{n-5}}{30405375} + \frac{3^{n-5} - (-3)^{n-5}}{256256000} + \frac{4^{n-6} + (-4)^{n-6}}{182432250} + \frac{3(6^{n-6} + (-6)^{n-6})}{11211200} + \frac{2^{n-8} + (-2)^{n-8}}{18243225} + \frac{2^{n-11} - (-2)^{n-11}}{6081075} + \frac{2^{n-7} - (-2)^{n-7} + 2^{n-12} + (-2)^{n-12}}{91216125},$$

$$\begin{aligned}
y_2(n, 9) = & \frac{(-1)^n + 1}{146313216000} + \frac{7^{n-2} + (-7)^{n-2}}{853991424000} + \frac{5^{n-2} + (-5)^{n-2}}{83691159552} + \frac{9^{n-4} + (-9)^{n-4}}{975822848000} \\
& + \frac{8^{n-5} - (-8)^{n-5}}{10854718875} + \frac{3^{n-5} - (-3)^{n-5}}{8712704000} + \frac{2(4^{n-5} - (-4)^{n-5})}{1550674125} \\
& + \frac{6^{n-6} + (-6)^{n-6}}{952952000} + \frac{3^{n-6} + (-3)^{n-6}}{871270400} + \frac{6^{n-7} - (-6)^{n-7}}{34034000} \\
& + \frac{2(4^{n-7} - (-4)^{n-7})}{1550674125} + \frac{3^{n-7} - (-3)^{n-7}}{536166400} + \frac{2^{n-7} - (-2)^{n-7}}{10854718875} \\
& + \frac{2^{n-8} + (-2)^{n-8}}{1550674125} + \frac{2^{n-11} - (-2)^{n-11}}{310134825} + \frac{2^{n-12} + (-2)^{n-12}}{1550674125}.
\end{aligned}$$

For  $n = 0$ , we have

$$y_2(0, 0) = 1, \quad y_2(0, 1) = 2, \quad y_2(0, 2) = \frac{2}{3}, \quad y_2(0, 3) = \frac{4}{45}, \quad y_2(0, 4) = \frac{2}{315},$$

and for  $k = 0, 1, 2, \dots, 9$  and  $n = 1, 2, \dots, 9$ , we use (20) to compute a few values of the numbers  $y_2(n, k)$ , as follows:

$n \setminus k$	0	1	2	3	4	5	6	7	8	9
1	0	0	0	0	0	0	0	0	0	0
2	0	1	$\frac{2}{3}$	$\frac{2}{15}$	$\frac{4}{315}$	$\frac{2}{2835}$	$\frac{4}{155925}$	$\frac{4}{6081075}$	$\frac{8}{638512875}$	$\frac{2}{10854718875}$
3	0	0	0	0	0	0	0	0	0	0
4	0	1	$\frac{5}{3}$	$\frac{8}{15}$	$\frac{22}{315}$	$\frac{2}{405}$	$\frac{34}{155925}$	$\frac{8}{1216215}$	$\frac{92}{638512875}$	$\frac{2}{834978375}$
5	0	0	0	0	0	0	0	0	0	0
6	0	1	$\frac{17}{3}$	$\frac{47}{15}$	$\frac{184}{315}$	$\frac{152}{2835}$	$\frac{454}{155925}$	$\frac{634}{6081075}$	$\frac{1688}{638512875}$	$\frac{542}{10854718875}$
7	0	0	0	0	0	0	0	0	0	0
8	0	1	$\frac{65}{3}$	$\frac{338}{15}$	$\frac{1957}{315}$	$\frac{2144}{2835}$	$\frac{7984}{155925}$	$\frac{2672}{1216215}$	$\frac{41462}{638512875}$	$\frac{15206}{10854718875}$
9	0	0	0	0	0	0	0	0	0	0

**Table 4:** Some numerical values of the numbers  $y_2(n, k)$ .

This function is related to the cosh  $t$ . That is,

$$F_{y_2}(t, k) = \frac{2^k}{(2k)!} (\cosh t + 1)^k.$$

By using this function, we get the following combinatorial sums:

**Theorem 12.** *Each of the following identities holds true:*

$$y_2(n, k; 1) = \frac{1}{(2k)!} \sum_{j=0}^k \binom{k}{j} 2^{k-j} \sum_{l=0}^j \binom{j}{l} (2l-j)^{2n}.$$

Also

$$\sum_{j=0}^k \binom{k}{j} 2^{k-j} \sum_{l=0}^j \binom{j}{l} (2l-j)^{2n+1} = 0.$$

*Proof.* By using (19), we have

$$\sum_{n=0}^{\infty} y_2(n, k) \frac{t^{2n}}{(2n)!} = \sum_{n=0}^{\infty} \left( \frac{1}{(2k)!} \sum_{j=0}^k \binom{k}{j} 2^{k-j} \sum_{l=0}^j \binom{j}{l} (2l-j)^{2n} \right) \frac{t^n}{n!}.$$

Comparing the coefficients of  $t^{2n}$  on both sides of the above equation, we obtain the desired result.  $\square$

By using (19), we obtain

$$F_{y_1}(t, 2k; 1)e^{-kt} = \frac{k!}{(2k)!} \sum_{v=0}^k F_{y_1}(t, v; 1)F_{y_1}(-t, k-v; 1).$$

By using the above functional equation, we obtain the following theorem:

**Theorem 13.** *The following identity holds:*

$$\sum_{j=0}^n \binom{n}{j} (-k)^{n-j} y_1(j, 2k; 1) = \frac{k!}{(2k)!} \sum_{j=0}^n (-1)^{n-j} \binom{n}{j} \sum_{v=0}^k y_1(j, v; 1) y_1(n-j, k-v; 1).$$

Recall that the following identity has very important applications in theory of double series and its applications ([29, Lemma 11, Eq-(7)]):

$$(21) \quad \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} A(n, k) = \sum_{n=0}^{\infty} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} A(n, n-2k),$$

where  $\lfloor x \rfloor$  denotes the greatest integer function.

**Theorem 14.** *The following identity holds:*

$$y_1(n, 2k; 1) = \sum_{j=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{2j} k^{n-2j} y_2(j, k; 1).$$

*Proof.* By using (19), we obtain the following functional equation:

$$F_{y_1}(t, 2k; 1) = F_{y_2}(t, k)e^{kt}.$$

Combining this equation with (8), we get

$$\sum_{n=0}^{\infty} y_1(n, 2k; 1) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \frac{(kt)^n}{n!} \sum_{n=0}^{\infty} y_2(n, k; 1) \frac{t^{2n}}{(2n)!}.$$

By using (21) in the above equation, we obtain

$$\sum_{n=0}^{\infty} y_1(n, 2k; 1) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left( \sum_{j=0}^{\lfloor \frac{n}{2} \rfloor} \frac{k^{n-2j}}{(2j)!(n-2j)!} y_2(j, k; 1) \right) t^n.$$

Comparing the coefficients of  $t^n$  on both sides of the above equation, we obtain the desired result.  $\square$

We now present a relation between the Lucas numbers  $L_n$  and the numbers  $y_2(n, k; 1)$  by the following theorem:

**Theorem 15.** *Let  $a + b = 1$ ,  $ab = -1$  and  $\frac{a-b}{2} = c = \frac{\sqrt{5}}{2}$ . Then*

$$L_n^{(k)} = \sum_{j=0}^k \binom{k}{j} (2j)! (-2)^{k-j} \sum_{m=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{2m} c^{2m} y_2(m, j; 1) \left(\frac{k}{2}\right)^{n-2m},$$

where  $L_n^{(k)}$  denote Lucas numbers of order  $k$ .

*Proof.* In [22, pp. 232-233] and [8], the Lucas numbers  $L_n$  are defined by means of the following generating function:

$$e^{at} + e^{bt} = \sum_{n=0}^{\infty} L_n \frac{t^n}{n!}.$$

From the above, we have

$$(22) \quad F_L(t, k; a, b) = (e^{at} + e^{bt})^k = \sum_{n=0}^{\infty} L_n^{(k)} \frac{t^n}{n!}.$$

where

$$L_n^{(k)} = \sum_{j=0}^n \binom{n}{j} L_n^{(m)} L_n^{(k-m)}.$$

By combining (22) with (16), we obtain the following functional equation

$$F_L(t, k; a, b) = e^{\frac{tk}{2}} \sum_{j=0}^k \binom{k}{j} (-2)^{k-j} (2j)! F_{y_2}(ct, j; 1).$$

Since  $F_{y_2}(ct, j; 1)$  is an even function, we have

$$\sum_{n=0}^{\infty} L_n^{(k)} \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left(\frac{k}{2}\right)^n \frac{t^n}{n!} \sum_{j=0}^k \binom{k}{j} (-2)^{k-j} (2j)! \sum_{m=0}^{\infty} y_2(m, j; 1) c^m \frac{t^{2m}}{(2m)!}.$$

Using (21) in the above equation, we get

$$\sum_{n=0}^{\infty} L_n^{(k)} \frac{t^n}{n!} = \sum_{n=0}^{\infty} \sum_{j=0}^k \binom{k}{j} (2j)! (-2)^{k-j} \sum_{m=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{2m} c^{2m} y_2(m, j; 1) \left(\frac{k}{2}\right)^{n-2m} \frac{t^n}{n!}.$$

Comparing the coefficients of  $t^n$  on both sides of the above equation, we obtain the desired result.  $\square$

We also present an identity including the Fibonacci numbers  $f_n$ , the Lucas numbers  $L_n$  and the numbers  $y_1(n, k; 1)$  by the following theorem:

**Theorem 16.** *Let  $a + b = 1$ ,  $ab = -1$  and  $\frac{a-b}{2} = c = \frac{\sqrt{5}}{2}$ . Then*

$$L_n^{(k)} = k! \sum_{j=0}^n \binom{n}{j} (2c)^{n-j} y_1(n-j, k; 1) ((a - 2ck^j) f_j + f_{j-1}).$$

*Proof.* We set

$$F_f(t, a, b) = \frac{e^{at} - e^{bt}}{a - b} = \sum_{n=0}^{\infty} f_n \frac{t^n}{n!}$$

(cf. [22, p. 232], [8]). By combining (22) and (8) with the above equation, we obtain the following functional equation

$$F_L(t, k; a, b) = k! F_{y_1}(2ct, k; 1) (e^{akt} - 2cF_f(kt, a, b)).$$

Therefore

$$\begin{aligned} \sum_{n=0}^{\infty} L_n^{(k)} \frac{t^n}{n!} &= k! \sum_{n=0}^{\infty} \sum_{j=0}^n \binom{n}{j} a^j (2c)^{n-j} y_1(n-j, k; 1) \frac{t^n}{n!} \\ &\quad - k! \sum_{n=0}^{\infty} \sum_{j=0}^n \binom{n}{j} (2c)^{n-j+1} y_1(n-j, k; 1) k^j f_j \frac{t^n}{n!}. \end{aligned}$$

After some elementary calculations and comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we obtain the desired result.  $\square$

#### 4.1. Recurrence relation for the numbers $y_2(n, k; \lambda)$

Here, taking derivative of (16), with respect to  $t$ , we give a recurrence relation for the numbers  $y_2(n, k; \lambda)$ .

**Theorem 17.** *Let  $k \in \mathbb{N}$ . Then*

$$y_2(n+1, k; \lambda) = ky_2(n, k; \lambda) - y_2(n, k-1; \lambda) - \lambda^{-1} \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} y_2(j, k-1; \lambda).$$

*Proof.* Taking derivative of (16), with respect to  $t$ , we obtain the following partial differential equation:

$$\frac{\partial}{\partial t} F_{y_2}(t, k; \lambda) = k F_{y_2}(t, k; \lambda) - F_{y_2}(t, k-1; \lambda) - \lambda^{-1} e^{-t} F_{y_2}(t, k-1; \lambda).$$

Combining (16) with the above equation, we obtain

$$\begin{aligned} \sum_{n=1}^{\infty} y_2(n, k; \lambda) \frac{t^{n-1}}{(n-1)!} &= k \sum_{n=0}^{\infty} y_2(n, k; \lambda) \frac{t^n}{n!} - \sum_{n=0}^{\infty} y_2(n, k-1; \lambda) \frac{t^n}{n!} \\ &\quad - \sum_{n=0}^{\infty} \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} y_2(j, k-1; \lambda) \frac{t^n}{n!}. \end{aligned}$$

After some elementary calculation, comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we obtain the desired result.  $\square$

**Theorem 18.** *Let  $n \in \mathbb{N}_0$  and  $k \in \mathbb{N}$ . Then*

$$2\lambda^2 \frac{\partial}{\partial \lambda} y_2(n, k; \lambda) = \lambda^2 \sum_{j=0}^n \binom{n}{j} y_2(j, k-1; \lambda) - \sum_{j=0}^n (-1)^{n-j} \binom{n}{j} y_2(j, k-1; \lambda).$$

*Proof.* Taking derivative of (16), with respect to  $\lambda$ , we obtain the following partial differential equation:

$$2\lambda^2 \frac{\partial}{\partial \lambda} F_{y_2}(t, k; \lambda) = F_{y_2}(t, k-1; \lambda) \left( e^t - \frac{1}{\lambda^2} e^{-t} \right).$$

Combining (16) with the above equation, we get

$$\begin{aligned} 2\lambda^2 \sum_{n=0}^{\infty} \frac{\partial}{\partial \lambda} y_2(n, k; \lambda) \frac{t^n}{n!} &= \sum_{n=0}^{\infty} \sum_{j=0}^n \binom{n}{j} y_2(j, k-1; \lambda) \frac{t^n}{n!} \\ &\quad - \frac{1}{\lambda^2} \sum_{n=0}^{\infty} \sum_{j=0}^n (-1)^{n-j} \binom{n}{j} y_2(j, k-1; \lambda) \frac{t^n}{n!}. \end{aligned}$$

After some elementary calculation, comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we get the desired result.  $\square$

## 5. $\lambda$ -central factorial numbers $C(n, k; \lambda)$

In this section, we define  $\lambda$ -central factorial numbers  $C(n, k; \lambda)$  by means of the following generating function: Let  $k \in \mathbb{N}_0$  and  $\lambda \in \mathbb{C}$ . Then

$$(23) \quad F_C(t, k; \lambda) = \frac{1}{(2k)!} (\lambda e^t + \lambda^{-1} e^{-t} - 2)^k = \sum_{n=0}^{\infty} C(n, k; \lambda) \frac{t^n}{n!}.$$

For  $\lambda = 1$ , we have the central factorial numbers

$$T(n, k) = C(n, k; 1)$$

(cf. [2], [7], [18], [35], [41]).

**Theorem 19.** *The following identity holds:*

$$C(n, k; \lambda^2) = 2^{-n}(2k)! \sum_{j=0}^n \binom{n}{j} C(j, k; \lambda) y_2(n-j, k; \lambda).$$

*Proof.* By using (16) and (23), we get the following functional equation:

$$F_C(2t, k; \lambda^2) = (2k)! F_C(t, k; \lambda) F_{y_2}(t, k; \lambda).$$

From this equation, we get

$$\sum_{n=0}^{\infty} C(n, k; \lambda^2) \frac{(2t)^n}{n!} = (2k)! \sum_{n=0}^{\infty} C(n, k; \lambda) \frac{t^n}{n!} \sum_{n=0}^{\infty} y_2(n, k; \lambda) \frac{t^n}{n!}.$$

Therefore

$$\sum_{n=0}^{\infty} C(n, k; \lambda^2) \frac{2^n t^n}{n!} = \sum_{n=0}^{\infty} (2k)! \sum_{j=0}^n \binom{n}{j} C(j, k; \lambda) y_2(n-j, k; \lambda) \frac{t^n}{n!}.$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we get the desired result.  $\square$

By using (7) and (19), we obtain the following functional equation:

$$F_T(t, k; 1) F_{y_2}(t, k; 1) = \frac{1}{(2k)!} F_T(2t, k; 1).$$

Combining the above equation with (7) and (19), we get

$$\frac{1}{(2k)!} \sum_{n=0}^{\infty} T(n, k) \frac{4^n t^{2n}}{(2n)!} = \sum_{n=0}^{\infty} y_2(n, k) \frac{t^n}{n!} \sum_{n=0}^{\infty} T(n, k) \frac{t^{2n}}{(2n)!}.$$

Therefore

$$\frac{1}{(2k)!} \sum_{n=0}^{\infty} 4^n T(n, k) \frac{t^{2n}}{(2n)!} = \sum_{n=0}^{\infty} \left( \sum_{j=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{2j} y_2(n-2j, k) T(j, k) \right) \frac{t^n}{n!}.$$

Comparing the coefficients of  $\frac{t^{2n}}{(2n)!}$  on both sides of the above equation, we obtain a relationship between the central factorial numbers  $T(n, k)$  and the numbers  $y_2(j, k)$  by the following theorem:

**Theorem 20.** *If  $n$  is even an integer, then we have*

$$T(n, k) = 4^{-n} (2k)! \sum_{j=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{2j} y_2(n-2j, k) T(j, k).$$

*If  $n$  is an odd integer, then we have*

$$\sum_{j=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{2j} y_2(n-2j, k) T(j, k) = 0.$$

**Remark 5.** *In [3], Alayont et al. have studied the rook polynomials, which count the number of ways of placing non-attacking rooks on a chess board. By using generalization of these polynomials, they gave the rook number interpretations of generalized central factorial and the Genocchi numbers.*

*In [2], Alayont and Krzywonos gave the following result for the classical central factorial numbers:*

*The number of ways to place  $k$  rooks on a size  $m$  triangle board in three dimensions is equal to  $T(m+1, m+1-k)$ , where  $0 \leq k \leq m$ .*

## 6. Application: in the binomial distribution and in the Bernstein polynomials

Let  $n$  be a nonnegative integer. For every function  $f : [0, 1] \rightarrow \mathbb{R}$  and the  $n^{\text{th}}$  Bernstein polynomial of  $f$  is defined by

$$B_n(f, x) = \sum_{k=0}^n \binom{n}{k} f\left(\frac{k}{n}\right) B_k^n(x),$$

where  $B_k^n(x)$  denotes the Bernstein basis functions:

$$B_k^n(x) = \binom{n}{k} x^k (1-x)^{n-k}$$

and  $x \in [0, 1]$ . Let  $(U_k)_{k \geq 1}$  be a sequence of independent distributed random variable having the uniform distribution on  $[0, 1]$  and defined by Adell *et al.* [1]:

$$S_n(x) = \sum_{k=1}^n 1_{[0, x]}(U_k).$$

In [1], it is well-know that,  $S_n(x)$  is a binomial random variable. That is the theory of Probability and Statistics, the binomial distribution is very useful. This distribution, with parameters  $n$  and probability  $x$ , is the discrete probability distribution. This distribution is defined as follows:

$$P(S_n(x) = k) = \binom{n}{k} x^k (1-x)^{n-k},$$



where  $k = 0, 1, 2, \dots, n$ . Let  $E$  denote mathematical expectation. Then

$$Ef\left(\frac{S_n(x)}{n}\right) = B_n(f, x)$$

(cf. [1]). For any  $x \in (0, 1)$ ,  $n \geq 2$ , and  $r > 1$ , Adel *et al.* [1] defined

$$(24) \quad E(S_n(x))^r = \sum_{k=0}^n \binom{n}{k} k^r x^k (1-x)^{n-k}.$$

Substituting  $x = \frac{1}{2}$  into (24), we get

$$(25) \quad E\left(S_n\left(\frac{1}{2}\right)\right)^r = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} k^r.$$

By combining (9) with (25), we arrive at the following theorem:

**Theorem 21.** *Let  $n \in \mathbb{N} \setminus \{1\}$ . Let  $r \in \mathbb{N}$ . Then*

$$y_1(r, n) = \frac{2^n}{n!} E\left(S_n\left(\frac{1}{2}\right)\right)^r.$$

Integrating (24) from 0 to 1, we get

$$\int_0^1 E(S_n(x))^r dx = \frac{1}{n+1} \sum_{k=0}^n k^r.$$

By substituting (2) into the above equation, after some elementary calculations, we get the following theorem:

**Theorem 22.** *The following identity holds:*

$$\int_0^1 E(S_n(x))^r dx = \frac{1}{(n+1)(r+1)} \sum_{j=0}^r \sum_{l=0}^{r+1-j} \binom{n+1}{j} \binom{r+1-j}{l} n^l B_l,$$

where  $B_l$  denotes the Bernoulli numbers.

## 7. Computation of the Euler numbers of negative order

In this section, we not only give elementary properties of the first and second kind Euler polynomials and numbers, but also compute the first kind of Apostol type Euler numbers associated with the numbers  $y_1(n, k; \lambda)$  and  $y_2(n, k; \lambda)$ .

The second kind Apostol type Euler polynomials of order  $k$ ,  $E_n^{*(k)}(x; \lambda)$  are given by means of the following generating function:

$$F_P(t, x; k, \lambda) = \left( \frac{2}{\lambda e^t + \lambda^{-1} e^{-t}} \right)^k e^{tx} = \sum_{n=0}^{\infty} E_n^{*(k)}(x; \lambda) \frac{t^n}{n!}.$$

Substituting  $x = 0$  into the above equation, we get the second kind Apostol type Euler numbers of order  $k$ , with  $k \geq 0$ ,  $E_n^{*(k)}(\lambda)$  by means of the following generating function:

$$F_N(t; k, \lambda) = \left( \frac{2}{\lambda e^t + \lambda^{-1} e^{-t}} \right)^k = \sum_{n=0}^{\infty} E_n^{*(k)}(\lambda) \frac{t^n}{n!}.$$

If we substitute  $k = \lambda = 1$  into the above generating function, then we have

$$E_n^* = E_n^{*(1)}(1).$$

Substituting  $x = 0$  into the equation (3) with  $-k$ , we get the first kind Apostol-Euler numbers of order  $-k$ ,  $E_n^{(-k)}(\lambda)$  are given by means of the following generating function:

$$(26) \quad G_E(t, -k; \lambda) = \left( \frac{\lambda e^t + 1}{2} \right)^k = \sum_{n=0}^{\infty} E_n^{(-k)}(\lambda) \frac{t^n}{n!}.$$

The second kind Apostol type Euler numbers of order  $-k$  are defined by means of the following generating function:

$$(27) \quad F_N(t; -k, \lambda) = \left( \frac{\lambda e^t + \lambda^{-1} e^{-t}}{2} \right)^k = \sum_{n=0}^{\infty} E_n^{*(-k)}(\lambda) \frac{t^n}{n!}.$$

The numbers  $E_n^{*(-k)}(\lambda)$  are related to the numbers  $E_n^{(-k)}(\lambda)$  and the Apostol Bernoulli numbers  $B_n^{(-k)}(\lambda)$  of the negative order. By using (27), we get the following functional equation:

$$F_N(t; -k, \lambda) = \sum_{j=0}^k \binom{k}{j} 2^{j-k} t^{k-j} G_E(t, -j; \lambda) H_B(-t, -k+j; \lambda^{-1}),$$

where

$$H_B(t, -k; \lambda) = \left( \frac{\lambda e^t - 1}{t} \right)^k = \sum_{n=0}^{\infty} B_n^{(-k)}(\lambda) \frac{t^n}{n!}$$

(cf. [25], [27], [39]). By using this equation, we get

$$\sum_{n=0}^{\infty} E_n^{*(-k)}(\lambda) \frac{t^n}{n!} = \sum_{j=0}^k \binom{k}{j} 2^{j-k} t^{k-j} \sum_{n=0}^{\infty} E_n^{(-j)}(\lambda) \frac{t^n}{n!} \sum_{n=0}^{\infty} B_n^{(-k+j)}(\lambda^{-1}) \frac{(-t)^n}{n!}.$$

Therefore

$$\begin{aligned} \sum_{n=0}^{\infty} E_n^{*(-k)}(\lambda) \frac{t^n}{n!} &= \sum_{n=0}^{\infty} \sum_{j=0}^k \binom{k}{j} \sum_{l=0}^{n-k+j} (-1)^{n+j-k-l} \binom{n-k+j}{l} \\ &\quad \times 2^{j-k} (n)_{k-j} E_l^{(-j)}(\lambda) B_{n+j-k-l}^{(-k+j)}(\lambda^{-1}) \frac{t^n}{n!}. \end{aligned}$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we arrive at the following theorem.

**Theorem 23.** *The following identity holds:*

$$\begin{aligned} E_n^{*(-k)}(\lambda) &= \sum_{j=0}^k \binom{k}{j} \sum_{l=0}^{n-k+j} (-1)^{n+j-k-l} \binom{n-k+j}{l} 2^{j-k} (n)_{k-j} \\ &\quad \times E_l^{(-j)}(\lambda) B_{n+j-k-l}^{(-k+j)}(\lambda^{-1}). \end{aligned}$$

We observe that the second kind Euler numbers of negative order  $E_n^{*(-k)}$  have been computed by Liu [24].

By using the numbers  $y_1(n, k; \lambda)$ , we are ready to compute the first kind Euler numbers of negative order.

Combining (8) and (26), we get

$$k!2^k \sum_{n=0}^{\infty} y_1(n, k; \lambda) \frac{t^n}{n!} = \sum_{n=0}^{\infty} E_n^{(-k)}(\lambda) \frac{t^n}{n!}.$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we arrive at the following theorem.

**Theorem 24.** *Let  $k \in \mathbb{N}_0$ . Then*

$$(28) \quad E_n^{(-k)}(\lambda) = k!2^{-k} y_1(n, k; \lambda).$$

**Remark 6.** *Substituting  $\lambda = 1$  into (28), we obtain the following explicit formula for the first kind Euler numbers of order  $-k$ :*

$$(29) \quad E_n^{(-k)} = 2^{-k} B(n, k).$$

From the equation (29), we see that

$$E_0^{(0)} = 1.$$

For  $k = 0, 1, 2, \dots, 7$  and  $n \in \mathbb{N}$ , we use (29) to compute a few values of the numbers

$E_n^{(-k)}$  as follows:

$$\begin{aligned}
 E_n^{(0)} &= 0, \\
 E_n^{(-1)} &= \frac{1}{2}, \\
 E_n^{(-2)} &= 2^{n-2} + \frac{1}{2}, \\
 E_n^{(-3)} &= \frac{3^n}{8} + 3 \cdot 2^{n-3} + \frac{3}{8}, \\
 E_n^{(-4)} &= \frac{3^n}{4} + 4^{n-2} + 3 \cdot 2^{n-3} + \frac{1}{4}, \\
 E_n^{(-5)} &= \frac{5^n}{32} + \frac{5 \cdot 3^n}{16} + \frac{5 \cdot 4^{n-2}}{2} + 5 \cdot 2^{n-4} + \frac{5}{32}, \\
 E_n^{(-6)} &= \frac{6^n}{64} + \frac{3 \cdot 5^n}{32} + \frac{5 \cdot 3^n}{16} + 15 \cdot 4^{n-3} + 15 \cdot 2^{n-6} + \frac{3}{32}, \\
 E_n^{(-7)} &= \frac{7^n}{128} + \frac{7 \cdot 6^n}{128} + \frac{21 \cdot 5^n}{128} + \frac{35 \cdot 3^n}{128} + \frac{35 \cdot 4^{n-3}}{2} + 21 \cdot 2^{n-7} + \frac{7}{128}, \\
 &\vdots
 \end{aligned}$$

That is for  $n = 0, 1, 2, \dots, 9$  and  $k = 0, 1, 2, \dots, 9$ , we compute a few values of the numbers  $E_n^{(-k)}$ , given by the above relations, as follows:

$n \setminus k$	0	1	2	3	4	5	6	7	8	9
0	1	1	1	1	1	1	1	1	1	1
1	0	$\frac{1}{2}$	1	$\frac{3}{2}$	2	$\frac{5}{2}$	3	$\frac{7}{2}$	4	$\frac{9}{2}$
2	0	$\frac{1}{2}$	$\frac{3}{2}$	3	5	$\frac{15}{2}$	$\frac{21}{2}$	14	18	$\frac{45}{2}$
3	0	$\frac{1}{2}$	$\frac{5}{2}$	$\frac{27}{4}$	14	25	$\frac{81}{2}$	$\frac{245}{4}$	88	$\frac{243}{2}$
4	0	$\frac{1}{2}$	$\frac{9}{2}$	$\frac{33}{2}$	$\frac{85}{2}$	90	168	287	459	$\frac{1395}{2}$
5	0	$\frac{1}{2}$	$\frac{17}{2}$	$\frac{171}{4}$	137	$\frac{1375}{4}$	738	1421	2524	4212
6	0	$\frac{1}{2}$	$\frac{33}{2}$	$\frac{231}{2}$	$\frac{925}{2}$	$\frac{5505}{4}$	$\frac{13587}{4}$	7364	14508	26550
7	0	$\frac{1}{2}$	$\frac{65}{2}$	$\frac{1287}{4}$	1619	5725	$\frac{65007}{4}$	$\frac{317275}{8}$	86608	173664
8	0	$\frac{1}{2}$	$\frac{129}{2}$	$\frac{1833}{2}$	$\frac{11665}{2}$	$\frac{49155}{2}$	$\frac{160671}{2}$	$\frac{441469}{2}$	$\frac{1068453}{2}$	1173240
9	0	$\frac{1}{2}$	$\frac{257}{2}$	$\frac{10611}{4}$	21497	$\frac{433225}{4}$	$\frac{816561}{2}$	$\frac{5055869}{4}$	3390874	$\frac{32620563}{4}$

**Table 5:** Some numerical values of the numbers  $E_n^{(-k)}$ .

**Theorem 25.** Let  $k \in \mathbb{N}_0$ . Then

$$y_2(n, k; \lambda) = \frac{2^k}{(2k)!} \sum_{l=0}^k \binom{k}{l} E_n^{*(-l)}(\lambda).$$

*Proof.* By using (16) and (27), we get the following functional equation:

$$F_{y_2}(t, k; \lambda) = \frac{2^k}{(2k)!} \sum_{l=0}^k \binom{k}{l} F_N(t; -l, \lambda).$$

From this equation, we obtain

$$\sum_{n=0}^{\infty} y_2(n, k; \lambda) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left( \frac{1}{(2k)!} \sum_{l=0}^k \binom{k}{l} 2^k E_n^{*(-l)}(\lambda) \right) \frac{t^n}{n!}.$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we obtain the desired result.  $\square$

**Theorem 26.** *Let  $k \in \mathbb{N}_0$ . Then*

$$y_2(n, k; \lambda) = \frac{2^k}{(2k)!} \sum_{l=0}^k \binom{k}{l} \lambda^{-l} E_n^{(-k)}(-l; \lambda).$$

*Proof.* By using (16) and (3), we get the following functional equation:

$$F_{y_2}(t, k; \lambda) = \frac{2^k}{(2k)!} \sum_{l=0}^k \binom{k}{l} \lambda^{-l} F_{P_1}(t, -l; -k, \lambda).$$

From this equation, we obtain

$$\sum_{n=0}^{\infty} y_2(n, k; \lambda) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left( \frac{2^k}{(2k)!} \sum_{l=0}^k \binom{k}{l} \lambda^{-l} E_n^{(-k)}(-l; \lambda) \right) \frac{t^n}{n!}.$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we obtain the desired result.  $\square$

By applying derivative operator to the generating function in (8), we give a relationship between the numbers  $y_1(n, k; \lambda)$  and  $E_n^{(-1)}(\lambda)$  as in the following theorem:

**Theorem 27.** *Let  $n, k \in \mathbb{N}_0$ . Then*

$$(30) \quad \begin{aligned} y_1(n+2, k; \lambda) &= k^2 y_1(n, k; \lambda) \\ &+ \frac{k(2k-3)}{2} \sum_{l=0}^n \binom{n}{l} y_1(n-l, k; \lambda) E_l(\lambda) \\ &+ \frac{k(k-1)}{4} \sum_{l=0}^n \binom{n}{l} E_l^{(2)}(\lambda) y_1(n-l, k; \lambda). \end{aligned}$$

*Let  $n \in \mathbb{N} \setminus \{1\}$ . Then*

$$(31) \quad y_1(n+2, k; \lambda) = k^2 y_1(n, k; \lambda) + y_1(n, k-2; \lambda) - (2k-1) y_1(n, k-1; \lambda).$$

*Proof of (30).* By applying derivative operator to (8) with respect to  $t$ , we obtain the following partial differential equation:

$$\begin{aligned} \frac{\partial^2}{\partial t^2} F_{y_1}(t, k; \lambda) &= k^2 F_{y_1}(t, k; \lambda) + \frac{k(2k-3)}{2} F_{P_1}(t, 0; 1, \lambda) F_{y_1}(t, k; \lambda) \\ &\quad + \frac{k(k-1)}{4} F_{P_1}(t, 0; 2, \lambda) F_{y_1}(t, k; \lambda). \end{aligned}$$

Combining (8) and (3) with the above equation, we get

$$\begin{aligned} &\sum_{n=2}^{\infty} y_1(n, k; \lambda) \frac{t^{n-2}}{(n-2)!} \\ &= k^2 \sum_{n=0}^{\infty} y_1(n, k; \lambda) \frac{t^n}{n!} + \frac{k(2k-3)}{2} \sum_{n=0}^{\infty} \sum_{l=0}^n \binom{n}{l} y_1(n-l, k; \lambda) E_l(\lambda) \frac{t^n}{n!} \\ &\quad + \frac{k(k-1)}{4} \sum_{n=0}^{\infty} \sum_{l=0}^n \binom{n}{l} E_l^{(2)}(\lambda) y_1(n-l, k; \lambda) \frac{t^n}{n!}. \end{aligned}$$

We make a suitable arrangement of the series and then compare the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, and we obtain the first assertion (30).  $\square$

*Proof of (31).* Similarly, by applying derivative operator to (8) with respect to  $t$ , we obtain the following partial differential equation:

$$\frac{\partial^2}{\partial t^2} F_{y_1}(t, k; \lambda) = k^2 F_{y_1}(t, k; \lambda) - (2k-1) F_{y_1}(t, k-1; \lambda) + F_{y_1}(t, k-2; \lambda).$$

Combining (8) with the above equation, we get

$$\begin{aligned} &\sum_{n=2}^{\infty} y_1(n, k; \lambda) \frac{t^{n-2}}{(n-2)!} \\ &= \sum_{n=0}^{\infty} (k^2 y_1(n, k; \lambda) - (2k-1) y_1(n, k-1; \lambda) + y_1(n, k-2; \lambda)) \frac{t^n}{n!}. \end{aligned}$$

Comparing the coefficients of  $\frac{t^n}{n!}$  on both sides of the above equation, we get the second assertion (31).  $\square$

## 8. Algorithms and Computation

In computer science and applied mathematics, one investigate information and computation and also their theoretical foundations. In these areas practical techniques are very important. Therefore algorithmic processes play a very important role in both areas. Thus, in this section, we give two algorithms for the computation of the numbers  $y_1(n, k; \lambda)$  and  $y_2(n, k; \lambda)$ .

---

**Algorithm 1** Let  $n$  be a positive integer and  $\lambda \neq 0$ . This algorithm will return value of the numbers  $y_1(n, k; \lambda)$  given by equation (9).

---

```

procedure  $y_1(n, k, \lambda)$ 
  Begin
  Inputs:
   $Y_1 \leftarrow 0$ 
  Outputs:
   $y_1(n, k, \lambda) \leftarrow Y_1$ 
  if  $n = 0$  and  $k = 0$  then
     $Y_1 = 1$ 
  else
    for all  $j$  in  $\{0, 1, 2, \dots, k\}$  do
       $Y_1 \leftarrow Y_1 + \text{Binomial.Coeff}(k, j) * \text{Power}(j, n) * \text{Power}(\lambda, j)$ 
    end for
  end if
   $Y_1 \leftarrow (1/k!) * Y_1$ 
  return  $Y_1$ 
end procedure

```

---



---

**Algorithm 2** Let  $n$  be a positive integer and  $\lambda \neq 0$ . This algorithm will return value of the numbers  $y_2(n, k; \lambda)$  given by equation (18) obtained by the  $y_1$  numbers in equation (9).

---

```

procedure  $y_2(n, k, \lambda)$ 
  Begin
  Inputs:
   $Y_2 \leftarrow 0$ 
  Outputs:
   $y_2(n, k, \lambda) \leftarrow Y_2$ 
  if  $n = 0$  and  $k = 0$  then
     $Y_2 = 1$ 
  else
    for all  $j$  in  $\{0, 1, 2, \dots, k\}$  do
      for all  $l$  in  $\{0, 1, 2, \dots, n\}$  do
         $Y_2 \leftarrow Y_2 + \text{Power}(-1, n - l) * \text{Binomial.Coeff}(n, l) * y_1(l, j, \lambda)$ 
         $* y_1(n - l, k - j, \text{Power}(\lambda, -1))$ 
      end for
    end for
  end if
   $Y_2 \leftarrow (k!/(2k!)) * Y_2$ 
  return  $Y_2$ 
end procedure

```

---

### 9. Combinatorial applications and further remarks

In this section, we discuss some combinatorial interpretations of these numbers, as well as the generalization of the central factorial numbers given in Section 3-5. These interpretations include the rook numbers and polynomials and combinatorial interpretation for the numbers  $y_1(n, k)$ . We see that our numbers are associated with known counting problems. By using counting techniques and generating function techniques, Bona [5] rederived several known properties and novel relations involving enumerative combinatorics and related areas. A very interesting further special case of the numbers  $B(n, k)$  is worthy of note by the work of Bona [5]. That is, in [5, P. 46, Exercise 3-4], Bona gave the following two exercises which are associated with the numbers  $B(n, k)$ :

**Exercise 3.** Find the number of ways to place  $n$  rooks on an  $n \times n$  chess board so that no two of them attack each other.

**Exercise 4.** How many ways are there to place some rooks on an  $n \times n$  chess board so that no two of them attack each other?

**Remark 7.** *Our numbers occur in combinatorics applications. In [5, p. 46, Exercise 3-4], Bona gave detailed and descriptive solution of these two exercises, which are related to the numbers  $B(n, k)$ , respectively, as follows:*

*There has to be one rook in each column. The first rook can be anywhere in its column ( $n$  possibilities). The second rook can be anywhere in its column except in the same row where the first rook is, which leaves  $n - 1$  possibilities. The third rook can be anywhere in its column, except in the rows taken by the first and second rook, which leaves  $n - 2$  possibilities, and so on, leading to  $n(n - 1) \cdots 2 \cdot 1 = n!$  possibilities.*

*Exercise 4. If we place  $k$  rooks, then we first need to choose the  $k$  columns in which these rooks will be placed. We can do that in  $\binom{n}{k}$  ways. Continuing the line of thought of the solution of the previous exercise, we can then place our  $k$  rooks into the chosen columns in  $(n)_k$  ways. Therefore, the total number of possibilities is*

$$\sum_{k=1}^n \binom{n}{k} (n)_k.$$

**Remark 8.** *In (14), for  $j < d$ , it is well-known that*

$$\binom{j}{d} = 0.$$

*Therefore, we arrive at solutions of Exercise 16 (a) in [5, p. 55, Exercise 16(a)] and also Exercise 10 [11, p. 126] as follows:*

$$2^{n-k} \binom{k}{d} = \sum_{j=d}^k \binom{k}{j} \binom{j}{d}.$$



## 10. Conclusions

In this paper, we have constructed some new families of special numbers with their generating functions. We give many properties of these numbers. These numbers are related to many well-known numbers, which are Bernoulli numbers, Euler numbers, Stirling numbers of the second kind, central factorial numbers and also related to the Golombek's problem [15] "Aufgabe 1088". We have discussed some combinatorial interpretations of these numbers. Besides, we give some applications about not only rook polynomials and numbers, but also combinatorial sum.

**Acknowledgment.** The present paper was supported by the *Scientific Research Project Administration of Akdeniz University*.

## REFERENCES

1. J. A. ADELL, J. BUSTAMANTE, J. M. QUESADA: *Estimates for the moments of Bernstein polynomials*. J. Math. Anal. Appl., **432** (2015), 114–128.
2. F. ALAYONT, N. KRZYWONOS: *Rook polynomials in three and higher dimensions*. Involve, **6**(1) (2013), 35–52.
3. F. ALAYONT, R. MOGER-REISCHER, R. SWIFT: *Rook number interpretations of generalized central factorial and Genocchi numbers*. [http://faculty.gvsu.edu/alayontf/notes/generalized\\_central\\_genocchi\\_numbers\\_preprint.pdf](http://faculty.gvsu.edu/alayontf/notes/generalized_central_genocchi_numbers_preprint.pdf).
4. A. BAYAD, Y. SIMSEK, H. M. SRIVASTAVA: *Some array type polynomials associated with special numbers and polynomials*. Appl. Math. Comput., **244** (2014), 149–157.
5. M. BONA: *Introduction to Enumerative Combinatorics*. The McGraw-Hill Companies, Inc. New York, 2007.
6. K. N. BOYADZHIEV: *Binomial transform and the backward difference*. arXiv preprint:1410.3014v2, <http://arxiv.org/abs/1410.3014v2>.
7. P. L. BUTZER, K. SCHMIDT, E. L. STARK, L. VOGT: *Central factorial numbers, their main properties and some applications*. Numer. Funct. Anal. Optim., **10**(5&6) (1989), 419–488.
8. P. F. BYRD: *New Relations between Fibonacci and Bernoulli Numbers*. Fibonacci Quart., **13** (1975), 111–114.
9. N. P. ČAKIĆ, G. V. MILOVANOVIC: *On generalized Stirling numbers and polynomials*. Math. Balkanica (N.S.), **18** (2004), 241–248.
10. C. H. CHANG, C.-W. HA: *A multiplication theorem for the Lerch zeta function and explicit representations of the Bernoulli and Euler polynomials*. J. Math. Anal. Appl., **315** (2006), 758–767.
11. C. A. CHARALAMBIDES: *Enumerative Combinatorics*. Chapman & Hall, Press Company, London, New York, 2002.
12. L. COMTET: *Advanced Combinatorics: The Art of Finite and Infinite Expansions*. Reidel, Dordrecht and Boston, 1974 (Translated from the French by J. W. Nienhuys).

13. G. B. DJORDJEVIC, G. V. MILOVANOVIC: *Special classes of polynomials*. University of Nis, Faculty of Technology Leskovac, 2014.
14. P. DOUBILET, G. C. ROTA, R. STANLEY: *On the foundations of combinatorial theory. VI. The idea of generating function*, Berkeley Symp. on Math. Statist. and Prob. Proc. Sixth Berkeley Symp. on Math. Statist. and Prob. Univ. of Calif. Press, **2** (1972), 267–318.
15. R. GOLOMBEK: *Aufgabe 1088*. *El. Math.*, **49** (1994), 126–127.
16. H. W. GOULD, J. QUAINANCE: *A Linear Binomial Recurrence and the Bell numbers and polynomials*. *Appl. Anal. Discrete Math.*, **1** (2007), 371–385.
17. Y. HE: *Some Results For Carlitz's  $q$ -Bernoulli Numbers And Polynomials*. *Appl. Anal. Discrete Math.*, **8** (2014), 304–319.
18. J. KANG, C. RYOO: *A research on the new polynomials involved with the central factorial numbers, Stirling numbers and others polynomials*. *J. Inequal. Appl.*, **26** (2014).
19. T. KIM:  *$q$ -Volkenborn integration*. *Russ. J. Math. Phys.*, **19** (2002), 288–299.
20. D. S. KIM, T. KIM: *Daehee numbers and polynomials*. *Appl. Math. Sci.*, **7** (2013), 5969–5976.
21. D. E. KNUTH: *The Art of Computer Programming. Volume 1 Fundamental Algorithms(Third Edition)*. Addison-Wesley, Massachusetts, 1997.
22. T. KOSHY: *Fibonacci and Lucas numbers with applications*. John Wiley & Sons, New York, 2001.
23. J. P. S. KUNG, G. C. ROTA, C. H. YAN: *Combinatorics: The Rota Way*. Cambridge University Press, New York, 2009.
24. G. LIU: *Generating functions and generalized Euler numbers*. *Proc. Japan Acad. Ser. A Math. Sci.*, **84** (2008), 29–34.
25. Q. M. LUO, H. M. SRIVASTAVA: *Some generalizations of the Apostol-Genocchi polynomials and the Stirling numbers of the second kind*. *Appl. Math. Comput.*, **217** (2011), 5702–5728.
26. H. OZDEN, Y. SIMSEK: *A new extension of  $q$ -Euler numbers and polynomials related to their interpolation functions*. *Appl. Math. Lett.*, **21**(9) (2008), 934–939.
27. H. OZDEN, Y. SIMSEK: *Modification and unification of the Apostol-type numbers and polynomials and their applications*. *Appl. Math. Comput.*, **235** (2014), 338–351.
28. H. OZDEN, Y. SIMSEK, H. M. SRIVASTAVA: *A unified presentation of the generating functions of the generalized Bernoulli, Euler and Genocchi polynomials*. *Comput. Math. Appl.*, **60** (2010), 2779–2787.
29. E. D. RAINVILLE: *Special functions*. The Macmillan Company, New York, 1960.
30. C. S. RYOO, T. KIM, L. C. JANG: *Some Relationships between the analogs of Euler numbers and polynomials*. *J. Inequal. Appl.*, **2007**(86052) (2007), 1–22.
31. Y. SIMSEK: *Special functions related to Dedekind-type DC-sums and their applications*. *Russ. J. Math. Phys.*, **17**(4) (2010), 495–508.
32. Y. SIMSEK: *On  $q$ -deformed Stirling numbers*. *Int. J. Math. Comput.*, **15** (2012), 70–80.
33. Y. SIMSEK: *Identities associated with generalized Stirling type numbers and Eulerian type polynomials*. *Math. Comput. Appl.*, **18** (2013), 251–263.

34. Y. SIMSEK: *Generating functions for generalized Stirling type numbers, array type polynomials, Eulerian type polynomials and their applications*. Fixed Point Theory Appl., **87** (2013), 343–355.
35. Y. SIMSEK: *Special numbers on analytic functions*. Applied Math. (Irvine), **5** (2014), 1091–1098.
36. Y. SIMSEK: *Analysis of the Bernstein basis functions: an approach to combinatorial sums involving binomial coefficients and Catalan numbers*. Math. Methods Appl. Sci., **38** (2015), 3007–3021.
37. M. R. SPIEGEL: *Calculus of finite differences and difference equations*. Schaum's Outline Series in Mathematics, McGraw-Hill Book Company, London, Toronto, 1971.
38. M. Z. SPIVEY: *Combinatorial Sums and Finite Differences*. Discrete Math., **307**(24) (2007), 3130–3146.
39. H. M. SRIVASTAVA: *Some generalizations and basic (or  $q$ -) extensions of the Bernoulli, Euler and Genocchi polynomials*. Appl. Math. Inf. Sci., **5** (2011), 390–444.
40. H. M. SRIVASTAVA, H. L. MANOCHA: *A Treatise on Generating Functions*. Ellis Horwood Limited Publisher, Chichester 1984.
41. H. M. SRIVASTAVA, G. D. LIU: *Some identities and congruences involving a certain family of numbers*. Russ. J. Math. Phys., **16** (2009), 536–542.
42. H. M. SRIVASTAVA, J. CHOI: *Zeta and  $q$ -Zeta Functions and Associated Series and Integrals*. Elsevier Science Publishers, Amsterdam, London and New York, 2012.
43. H. M. SRIVASTAVA, T. KIM, Y. SIMSEK:  *$q$ -Bernoulli numbers and polynomials associated with multiple  $q$ -zeta functions and basic  $L$ -series*. Russ. J. Math. Phys., **12** (2005), 241–268.
44. H. M. SRIVASTAVA, M. A. OZARSLAN, C. KAANOGLU: *Some families of generating functions for a certain class of three-variable polynomials*. Integral Transforms Spec. Funct., **21** (2010), 885–896.
45. C. F. WEI, F. QI: *Several closed expressions for the Euler Numbers*. J. Inequal. Appl., **2015**(219) (2015), 1–8.

**Yilmaz Simsek**

Department of Mathematics,

Faculty of Science

University of Akdeniz

TR-07058 Antalya,

Turkey.

E-mail: [ysimsek@akdeniz.edu.tr](mailto:ysimsek@akdeniz.edu.tr)

(Received 29.06.2016)

(Revised 08.03.2017)