

A DIAGONAL RECURRENCE RELATION FOR THE STIRLING NUMBERS OF THE FIRST KIND

Feng Qi and Bai-Ni Guo*

In the paper, the authors present an explicit form for a family of inhomogeneous linear ordinary differential equations, find a more significant expression for all derivatives of a function related to the solution to the family of inhomogeneous linear ordinary differential equations in terms of the Lerch transcendent, establish an explicit formula for computing all derivatives of the solution to the family of inhomogeneous linear ordinary differential equations, acquire the absolute monotonicity, complete monotonicity, the Bernstein function property of several functions related to the solution to the family of inhomogeneous linear ordinary differential equations, discover a diagonal recurrence relation of the Stirling numbers of the first kind, and derive an inequality for bounding the logarithmic function.

1. MAIN RESULTS

In [7, Section 3, Theorem 1], by inductive argument, it was proved that the family of inhomogeneous linear ordinary differential equations

$$(1) \quad (\sqrt{1-4t} - 1)F^{(n)}(t) + \sum_{i=1}^n \frac{a_{n,i}}{(1-4t)^{n-i+1/2}} F^{(i-1)}(t) = \frac{a_{n,0}}{(1-4t)^n}$$

for $n \in \mathbb{N}$ have a solution

$$(2) \quad F(t) = \begin{cases} \frac{1}{2} \frac{\ln(1-4t)}{\sqrt{1-4t} - 1}, & 0 \neq t < \frac{1}{4}; \\ 1, & t = 0, \end{cases}$$

*Corresponding author. Feng Qi

2010 Mathematics Subject Classification. 11B73, 11M35, 26A48.

Keywords and Phrases. diagonal recurrence relation; inhomogeneous linear ordinary differential equation; Stirling numbers of the first kind; Lerch transcendent; complete monotonicity.

where

$$\begin{aligned} a_{n,0} &= -2^{2n-1}(n-1)!, & a_{n,1} &= -2^n(2n-3)!!, & a_{n,n} &= -2n, \\ a_{n,i} &= -S_{n-i+2,i-2}2^{n-i+1}(2n-2i-1)!!, & 2 \leq i \leq n-1, \\ S_{n,0} &= n, & S_{n,j} &= \sum_{\ell=1}^n S_{\ell,j-1}, & j \geq 1. \end{aligned}$$

This is a core conclusion in the paper [7].

It is obvious that the above sequence $a_{n,i}$ for $n \in \mathbb{N}$ and $0 \leq i \leq n$ is given recursively. It is natural to ask for a question: can one find an explicit expression of the above sequence $a_{n,i}$ for $n \in \mathbb{N}$ and $0 \leq i \leq n$? In other words, can one find an explicit form for the family of inhomogeneous linear ordinary differential equations (1)? Our answer to this question can be stated as Theorem 1 below.

Theorem 1. For $n \in \mathbb{N}$, the family of inhomogeneous linear ordinary differential equations

$$\begin{aligned} (1 - \sqrt{1-4t})(1-4t)^n F^{(n)}(t) + 2n(1-4t)^{n-1/2} F^{(n-1)}(t) \\ + 2^n \sum_{r=0}^{n-2} \binom{n}{r} \frac{(2n-2r+1)!!}{2^r} (1-4t)^{r+1/2} F^{(r)}(t) = 2^{2n-1}(n-1)! \end{aligned}$$

has the solution (2).

We observe that the function $F(t)$ is a composite of the functions

$$f(x) = \begin{cases} \frac{\ln x}{x-1}, & 0 < x \neq 1 \\ 1, & x = 1 \end{cases}$$

and $x = h(t) = \sqrt{1-4t}$. By the Leibniz theorem for differentiation of a product, it is not difficult to obtain

$$(3) \quad f^{(n)}(x) = \frac{(-1)^n n!}{(x-1)^{n+1}} \left[\ln x - \sum_{k=1}^n \frac{1}{k} \left(\frac{x-1}{x} \right)^k \right], \quad n \geq 0.$$

Recall from [3, Chapter 14], [8, Chapter XIII], [28, Chapter 1], and [30, Chapter IV] that a function f is said to be completely monotonic on an interval $I \subseteq \mathbb{R}$ if f has derivatives of all orders on I and $(-1)^n f^{(n)}(t) \geq 0$ for all $t \in I$ and $n \in \{0\} \cup \mathbb{N}$. Recall from [22, p. 1161] and [28, Chapter 3] that a function $f : I \subseteq (-\infty, \infty) \rightarrow [0, \infty)$ is called a Bernstein function on I if $f(t)$ has derivatives of all orders and $f'(t)$ is completely monotonic on I .

The second main result of this paper is to present a more significant expression than (3) for the n th derivative of the function $f(x)$ in terms of the Lerch transcendent

$$\Phi(z, s, a) = \sum_{k=0}^{\infty} \frac{z^k}{(a+k)^s}, \quad a \neq 0, -1, \dots$$

From the more significant expression, we obtain complete monotonicity of the functions $f(x)$ and $x^{n+1}f^{(n)}(x)$ and derive an inequality of the logarithmic function $\ln x$.

Theorem 2. *For $n \geq 0$, the n th derivative of the function $f(x)$ can be expressed by*

$$(4) \quad f^{(n)}(x) = \begin{cases} (-1)^n \frac{n!}{x^{n+1}} \Phi\left(\frac{x-1}{x}, 1, n+1\right), & 0 < x \neq 1; \\ (-1)^n \frac{n!}{n+1}, & x = 1. \end{cases}$$

Consequently,

1. the function $f(x)$ is completely monotonic on $(0, \infty)$;
2. the functions $x^{n+1}f^{(n)}(x)$ for all $n \geq 0$ are Bernstein functions on $(0, \infty)$;
3. the inequality

$$(5) \quad \ln x > \sum_{k=1}^n \frac{1}{k} \left(\frac{x-1}{x}\right)^k, \quad n \in \mathbb{N}$$

(a) holds

- i. either for $x > 1$ and all $n \in \mathbb{N}$,
- ii. or for $0 < x < 1$ and odd n ;

(b) reverses for $0 < x < 1$ and even n .

Recall from [5] and [30, Chapter IV] that a function f is said to be absolutely monotonic on an interval I if it has derivatives of all orders and $f^{(k-1)}(t) \geq 0$ for $t \in I$ and $k \in \mathbb{N}$.

The third main result of this paper is an explicit formula for the n th derivative of $F(t)$. From the explicit formula, we deduce the absolute monotonicity of the function $F(t)$.

Theorem 3. *The n th derivative of the function $F(t)$ can be expressed by*

$$F^{(n)}(t) = \frac{2^n}{(1-4t)^{n+1/2}} \sum_{\ell=0}^n \ell! [2(n-\ell)-1]!! \binom{2n-\ell-1}{\ell-1} \Phi\left(\frac{\sqrt{1-4t}-1}{\sqrt{1-4t}}, 1, \ell+1\right).$$

Consequently, the function $F(t)$ is absolutely monotonic on $(-\infty, \frac{1}{4})$ and, equivalently, the function $F(-t)$ is completely monotonic on $(-\frac{1}{4}, \infty)$.

In combinatorics [4, 11, 13], the Stirling number of the first kind $s(n, k)$ can be defined such that the number of permutations of n elements which contain exactly k permutation cycles is the nonnegative number $|s(n, k)| = (-1)^{n-k} s(n, k)$.

In [11, 12], diagonal recurrence relations for the Stirling numbers of the first and second kinds were discovered.

The fourth main result of this paper is a diagonal recurrence relation of the Stirling numbers of the first kind $s(n, k)$.

Theorem 4. *The Stirling numbers of the first kind $s(n, k)$ satisfy the diagonal recurrence relation*

$$(6) \quad \frac{s(n+k, k)}{\binom{n+k}{k}} = \sum_{\ell=0}^n (-1)^\ell \frac{\langle k \rangle_\ell}{\ell!} \sum_{m=1}^{\ell} (-1)^m \binom{\ell}{m} \frac{s(n+m, m)}{\binom{n+m}{m}},$$

where

$$\langle x \rangle_n = \prod_{k=0}^{n-1} (x-k) = \begin{cases} x(x-1)\cdots(x-n+1), & n \geq 1 \\ 1, & n = 0 \end{cases}$$

is the falling factorial.

2. LEMMAS

In order to prove our main results, we recall several lemmas below.

Lemma 1 ([4, p. 134, Theorem A] and [4, p. 139, Theorem C]). *For $n \geq k \geq 0$, the Bell polynomials of the second kind, or say, partial Bell polynomials, denoted by $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$, are defined by*

$$B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{1 \leq i \leq n, \ell_i \in \{0\} \cup \mathbb{N} \\ \sum_{i=1}^n i \ell_i = n \\ \sum_{i=1}^n \ell_i = k}} \frac{n!}{\prod_{i=1}^{n-k+1} \ell_i!} \prod_{i=1}^{n-k+1} \left(\frac{x_i}{i!}\right)^{\ell_i}.$$

The Faà di Bruno formula can be described in terms of the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ by

$$(7) \quad \frac{d^n}{dt^n} f \circ h(t) = \sum_{k=0}^n f^{(k)}(h(t)) B_{n,k}(h'(t), h''(t), \dots, h^{(n-k+1)}(t)).$$

The Faà di Bruno formula (7) and the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ are useful tools in combinatorics and mathematical analysis and they were often used to compute the higher-order derivatives for a composite function. For example, see [31, 32].

Lemma 2 ([4, p. 135]). *For complex numbers a and b , we have*

$$(8) \quad B_{n,k}(abx_1, ab^2x_2, \dots, ab^{n-k+1}x_{n-k+1}) = a^k b^n B_{n,k}(x_1, x_2, \dots, x_{n-k+1}).$$

Lemma 3 ([4, p. 135, Theorem B] and [11, Theorem 1.1]). *For $n \geq k \geq 0$, we have*

$$(9) \quad B_{n,k}(1!, 2!, \dots, (n-k+1)!) = \binom{n-1}{k-1} \frac{n!}{k!}$$

and

$$(10) \quad B_{n,k}\left(\frac{1!}{2}, \frac{2!}{3}, \dots, \frac{(n-k+1)!}{n-k+2}\right) = (-1)^{n-k} \frac{1}{k!} \sum_{m=0}^k (-1)^m \binom{k}{m} \frac{s(n+m, m)}{\binom{n+m}{m}}.$$

Lemma 4 ([1, p. 40]). *Let $p = p(x)$ and $q = q(x) \neq 0$ be two differentiable functions. Then*

$$(11) \quad \left[\frac{p(x)}{q(x)} \right]^{(k)} = \frac{(-1)^k}{q^{k+1}} \begin{vmatrix} p & q & 0 & \cdots & 0 & 0 \\ p' & q' & q & \cdots & 0 & 0 \\ p'' & q'' & \binom{2}{1}q' & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ p^{(k-2)} & q^{(k-2)} & \binom{k-2}{1}q^{(k-3)} & \cdots & q & 0 \\ p^{(k-1)} & q^{(k-1)} & \binom{k-1}{1}q^{(k-2)} & \cdots & \binom{k-1}{k-2}q' & q \\ p^{(k)} & q^{(k)} & \binom{k}{1}q^{(k-1)} & \cdots & \binom{k}{k-2}q'' & \binom{k}{k-1}q' \end{vmatrix}$$

for $k \geq 0$. In other words, the formula (11) can be rewritten as

$$(12) \quad \frac{d^k}{dx^k} \left[\frac{p(x)}{q(x)} \right] = \frac{(-1)^k}{q^{k+1}(x)} |W_{(k+1) \times (k+1)}(x)|,$$

where $|W_{(k+1) \times (k+1)}(x)|$ denotes the determinant of the $(k+1) \times (k+1)$ matrix

$$W_{(k+1) \times (k+1)}(x) = (U_{(k+1) \times 1}(x) \quad V_{(k+1) \times k}(x)),$$

the quantity $U_{(k+1) \times 1}(x)$ is a $(k+1) \times 1$ matrix whose elements $u_{\ell,1}(x) = p^{(\ell-1)}(x)$ for $1 \leq \ell \leq k+1$, and $V_{(k+1) \times k}(x)$ is a $(k+1) \times k$ matrix whose elements

$$v_{i,j}(x) = \begin{cases} \binom{i-1}{j-1} q^{(i-j)}(x), & i-j \geq 0 \\ 0, & i-j < 0 \end{cases}$$

for $1 \leq i \leq k+1$ and $1 \leq j \leq k$.

Remark 1. The formula (12) in Lemma 4 has been applied in the papers [6, 10, 14, 15, 16, 17, 18, 19, 21, 23, 24, 25, 26, 27, 29] to express the Apostol–Bernoulli polynomials, the Cauchy product of central Delannoy numbers, the Bernoulli polynomials, the Schröder numbers, the (generalized) Fibonacci polynomials, the Catalan numbers, derangement numbers, and the Euler numbers and polynomials in terms of the Hessenberg and tridiagonal determinants. This implies that Lemma 4 is an effectual tool to express some mathematical quantities in terms of the Hessenberg and tridiagonal determinants.

Lemma 5 ([2, p. 222, Theorem] and [27, Remark 3]). *Let $M_0 = 1$ and*

$$M_n = \begin{vmatrix} m_{1,1} & m_{1,2} & 0 & \cdots & 0 & 0 \\ m_{2,1} & m_{2,2} & m_{2,3} & \cdots & 0 & 0 \\ m_{3,1} & m_{3,2} & m_{3,3} & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ m_{n-2,1} & m_{n-2,2} & m_{n-2,3} & \cdots & m_{n-2,n-1} & 0 \\ m_{n-1,1} & m_{n-1,2} & m_{n-1,3} & \cdots & m_{n-1,n-1} & m_{n-1,n} \\ m_{n,1} & m_{n,2} & m_{n,3} & \cdots & m_{n,n-1} & m_{n,n} \end{vmatrix}$$

for $n \in \mathbb{N}$. Then the sequence M_n for $n \geq 0$ satisfies $M_1 = m_{1,1}$ and

$$(13) \quad M_n = m_{n,n}M_{n-1} + \sum_{r=1}^{n-1} (-1)^{n-r} m_{n,r} \left(\prod_{j=r}^{n-1} m_{j,j+1} \right) M_{r-1}, \quad n \geq 2.$$

Lemma 6. *The Lerch transcendent $\Phi(z, s, a)$ satisfies*

$$(14) \quad \Phi(z, s, a) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{x^{s-1} e^{-ax}}{1 - ze^{-x}} dx, \quad \Re(s), \Re(a) > 0, \quad z \in \mathbb{C} \setminus [1, \infty)$$

and

$$(15) \quad \Phi(t, 1, n + 1) = -\frac{\ln(1 - t)}{t^{n+1}} - \sum_{k=1}^n \frac{1}{n - k + 1} \frac{1}{t^k}, \quad n \geq 0,$$

where

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt, \quad \Re(z) > 0$$

denotes the classical Euler gamma function. Consequently, the Lerch transcendent $\Phi(x, s, a)$ for $s, a > 0$ is an absolutely monotonic function of $x \in (-\infty, 1)$ and the function $\Phi\left(\frac{x-1}{x}, 1, n + 1\right)$ is a Bernstein function of $x \in (0, \infty)$.

Proof. The integral representation (14) can be found in [9, p. 612, Entry 25.14.5].

From (14), it follows that

$$\frac{\partial^n}{\partial z^n} \Phi(z, s, a) = \frac{n!}{\Gamma(s)} \int_0^\infty \frac{x^{s-1} e^{-(a+n)x}}{(1 - ze^{-x})^{n+1}} dx$$

for $\Re(s), \Re(a) > 0$ and $z \in \mathbb{C} \setminus [1, \infty)$. Hence, the Lerch transcendent $\Phi(x, s, a)$ for $s, a > 0$ is an absolutely monotonic function of $x \in (-\infty, 1)$.

Since

$$\ln(1 - t) = -\sum_{k=1}^\infty \frac{t^k}{k}, \quad |t| < 1,$$

we have

$$\begin{aligned} & -\frac{\ln(1 - t)}{t^{n+1}} - \sum_{k=1}^n \frac{1}{n - k + 1} \frac{1}{t^k} = \sum_{k=1}^\infty \frac{1}{k} \frac{1}{t^{n-k+1}} - \sum_{\ell=1}^n \frac{1}{\ell} \frac{1}{t^{n-\ell+1}} \\ & = \sum_{k=n+1}^\infty \frac{1}{k} \frac{1}{t^{n-k+1}} = \sum_{k=0}^\infty \frac{1}{n + k + 1} \frac{1}{t^k} = \Phi(t, 1, n + 1). \end{aligned}$$

The identity (15) is thus proved.

A direct computation by employing (7), (8), and (9) gives

$$\frac{\partial^k}{\partial x^k} \Phi\left(\frac{x-1}{x}, 1, n + 1\right) = \sum_{\ell=0}^k \frac{\partial^\ell}{\partial u^\ell} \Phi(u, 1, n + 1)$$

$$\begin{aligned}
 & \times B_{k,\ell} \left(\frac{1!}{x^2}, -\frac{2!}{x^3}, \dots, (-1)^{k-\ell} \frac{(k-\ell+1)!}{x^{k-\ell+2}} \right) \\
 &= \sum_{\ell=0}^k \frac{\partial^\ell}{\partial u^\ell} \Phi(u, 1, n+1) \frac{(-1)^{k+\ell}}{x^{k+\ell}} B_{k,\ell}(1!, 2!, \dots, (k-\ell+1)!) \\
 &= \frac{(-1)^k}{x^k} \sum_{\ell=0}^k \frac{\ell!}{\Gamma(1)} \int_0^\infty \frac{e^{-(n+\ell+1)t}}{(1-ue^{-t})^{\ell+1}} dt \frac{(-1)^\ell}{x^\ell} \binom{k-1}{\ell-1} \frac{k!}{\ell!} \\
 &= \frac{(-1)^k k!}{x^k} \int_0^\infty e^{-(n+1)t} \sum_{\ell=0}^k \frac{e^{-\ell t}}{(1-ue^{-t})^{\ell+1}} \frac{(-1)^\ell}{x^\ell} \binom{k-1}{\ell-1} dt \\
 &= \frac{(-1)^{k+1} k!}{x^k} \int_0^\infty e^{-(n+1)t} \frac{e^t}{(e^t-1)(e^t x - x + 1)} \left[\frac{(e^t-1)x}{(e^t-1)x+1} \right]^k dt \\
 &= (-1)^{k+1} k! \int_0^\infty \frac{(e^t-1)^{k-1}}{[(e^t-1)x+1]^{k+1} e^{nt}} dt
 \end{aligned}$$

for $k \in \mathbb{N}$, where $u = 1 - \frac{1}{x}$. Thus, the function $\Phi(\frac{x-1}{x}, 1, n+1)$ is a Bernstein function on $(0, \infty)$. The proof of Lemma 6 is complete. \square

Lemma 7 ([25, Theorem 4]). *Let $h_{a,b}(x) = \sqrt{a+bx}$ for $a, b \in \mathbb{R}$ and $b \neq 0$ and let $n \in \mathbb{N}$. Then the Bell polynomials of the second kind $B_{n,k}$ satisfy*

$$\begin{aligned}
 (16) \quad & B_{n,k}(h'_{a,b}(x), h''_{a,b}(x), \dots, h_{a,b}^{(n-k+1)}(x)) \\
 &= (-1)^{n+k} [2(n-k)-1]!! \left(\frac{b}{2}\right)^n \binom{2n-k-1}{2(n-k)} \frac{1}{(a+bx)^{n-k/2}}.
 \end{aligned}$$

3. PROOFS OF MAIN RESULTS

We now start out to prove our main results.

Proof of Theorem 1. By virtue of Lemma 4, we have

$$\begin{aligned}
 2F^{(n)}(t) &= \left[\frac{\ln(1-4t)}{\sqrt{1-4t}-1} \right]^{(n)} = \frac{(-1)^n}{(\sqrt{1-4t}-1)^{n+1}} M_{n+1} \triangleq \frac{(-1)^n}{(\sqrt{1-4t}-1)^{n+1}} \\
 &\times \begin{vmatrix} \ln(1-4t) & \sqrt{1-4t}-1 & 0 & \cdots & 0 \\ \frac{-4}{1-4t} & \frac{-2}{(1-4t)^{1/2}} & \sqrt{1-4t}-1 & \cdots & 0 \\ \frac{-4^2 1!}{(1-4t)^2} & \frac{-2^2 1!!}{(1-4t)^{3/2}} & \binom{2}{1} \frac{-2}{(1-4t)^{1/2}} & \cdots & 0 \\ \frac{-4^3 2!}{(1-4t)^3} & \frac{-2^3 3!!}{(1-4t)^{5/2}} & \binom{3}{1} \frac{-2^2 1!!}{(1-4t)^{3/2}} & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \frac{-4^{n-2} (n-3)!}{(1-4t)^{n-2}} & \frac{-2^{n-2} (2n-7)!!}{(1-4t)^{(2n-5)/2}} & \binom{n-2}{1} \frac{-2^{n-3} (2n-9)!!}{(1-4t)^{(2n-7)/2}} & \cdots & 0 \\ \frac{-4^{n-1} (n-2)!}{(1-4t)^{n-1}} & \frac{-2^{n-1} (2n-5)!!}{(1-4t)^{(2n-3)/2}} & \binom{n-1}{1} \frac{-2^{n-2} (2n-7)!!}{(1-4t)^{(2n-5)/2}} & \cdots & \sqrt{1-4t}-1 \\ \frac{-4^n (n-1)!}{(1-4t)^n} & \frac{-2^n (2n-3)!!}{(1-4t)^{(2n-1)/2}} & \binom{n}{1} \frac{-2^{n-1} (2n-5)!!}{(1-4t)^{(2n-3)/2}} & \cdots & \binom{n}{n-1} \frac{-2}{(1-4t)^{1/2}} \end{vmatrix}.
 \end{aligned}$$

Further applying the recurrence relation (13) to the above determinant yields

$$M_{n+1} = \binom{n}{n-1} \frac{-2}{(1-4t)^{1/2}} M_n + (-1)^n \frac{-4^n(n-1)!}{(1-4t)^n} (\sqrt{1-4t}-1)^n$$

$$+ \sum_{r=2}^n (-1)^{n-r+1} \binom{n}{r-2} \frac{-2^{n-r+2}[2(n-r)+1]!!}{(1-4t)^{[2(n-r)+3]/2}} (\sqrt{1-4t}-1)^{n-r+1} M_{r-1}$$

for $n \in \mathbb{N}$. This equality can be rewritten as

$$\frac{(-1)^n}{(\sqrt{1-4t}-1)^{n+1}} M_{n+1} = \frac{1}{\sqrt{1-4t}-1} \left[\binom{n}{n-1} \frac{2}{(1-4t)^{1/2}} \frac{(-1)^{n-1}}{(\sqrt{1-4t}-1)^n} M_n \right.$$

$$\left. - \frac{4^n(n-1)!}{(1-4t)^n} + \sum_{r=2}^n \binom{n}{r-2} \frac{2^{n-r+2}[2(n-r)+1]!!}{(1-4t)^{[2(n-r)+3]/2}} \frac{(-1)^{r-2}}{(\sqrt{1-4t}-1)^{r-1}} M_{r-1} \right]$$

for $n \in \mathbb{N}$. In a word, we obtain

$$2F^{(n)}(t) = \frac{1}{\sqrt{1-4t}-1} \left[\binom{n}{n-1} \frac{2}{(1-4t)^{1/2}} 2F^{(n-1)}(t) \right.$$

$$\left. + \sum_{r=2}^n \binom{n}{r-2} \frac{2^{n-r+2}[2(n-r)+1]!!}{(1-4t)^{[2(n-r)+3]/2}} 2F^{(r-2)}(t) - \frac{4^n(n-1)!}{(1-4t)^n} \right], \quad n \in \mathbb{N}.$$

Theorem 1 is thus proved. □

Proof of Theorem 2. Applying $p(x) = \ln x$ and $q(x) = x-1$ to Lemma 4, expanding the obtained determinant according to the last row consecutively, and making use of (15) yield

$$\left(\frac{\ln x}{x-1} \right)^{(n)} = \frac{(-1)^n}{(x-1)^{n+1}}$$

$$\times \begin{vmatrix} \ln x & x-1 & 0 & 0 & \cdots & 0 & 0 \\ \frac{1}{x} & 1 & x-1 & 0 & \cdots & 0 & 0 \\ -\frac{1}{x^2} & 0 & 2 & x-1 & \cdots & 0 & 0 \\ \frac{2}{x^3} & 0 & 0 & 3 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ (-1)^{n-2} \frac{(n-2)!}{x^{n-1}} & 0 & 0 & 0 & \cdots & n-1 & x-1 \\ (-1)^{n-1} \frac{(n-1)!}{x^n} & 0 & 0 & 0 & \cdots & 0 & n \end{vmatrix}$$

$$= \frac{(-1)^n n}{(x-1)^{n+1}} \begin{vmatrix} \ln x & x-1 & 0 & 0 & \cdots & 0 & 0 \\ \frac{1}{x} & 1 & x-1 & 0 & \cdots & 0 & 0 \\ -\frac{1}{x^2} & 0 & 2 & x-1 & \cdots & 0 & 0 \\ \frac{2}{x^3} & 0 & 0 & 3 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ (-1)^{n-3} \frac{(n-3)!}{x^{n-2}} & 0 & 0 & 0 & \cdots & n-2 & x-1 \\ (-1)^{n-2} \frac{(n-2)!}{x^{n-1}} & 0 & 0 & 0 & \cdots & 0 & n-1 \end{vmatrix}$$

$$\begin{aligned}
 &+ (-1)^{n-1} \frac{(n-1)!}{x^n(x-1)} \\
 = &\frac{(-1)^n n(n-1)}{(x-1)^{n+1}} \begin{vmatrix} \ln x & x-1 & 0 & 0 & \cdots & 0 & 0 \\ \frac{1}{x} & 1 & x-1 & 0 & \cdots & 0 & 0 \\ -\frac{1}{x^2} & 0 & 2 & x-1 & \cdots & 0 & 0 \\ \frac{2}{x^3} & 0 & 0 & 3 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ (-1)^{n-4} \frac{(n-4)!}{x^{n-3}} & 0 & 0 & 0 & \cdots & n-3 & x-1 \\ (-1)^{n-3} \frac{(n-3)!}{x^{n-2}} & 0 & 0 & 0 & \cdots & 0 & n-2 \end{vmatrix} \\
 &+ (-1)^{n-1} \frac{(n-2)!n}{x^{n-1}(x-1)^2} + (-1)^{n-1} \frac{(n-1)!}{x^n(x-1)} \\
 = &\frac{(-1)^n \langle n \rangle_{n-1}}{(x-1)^{n+1}} \begin{vmatrix} \ln x & x-1 \\ \frac{1}{x} & 1 \end{vmatrix} + (-1)^{n-1} \sum_{k=1}^{n-1} \frac{(n-k)! \langle n \rangle_{k-1}}{x^{n-k+1}(x-1)^k} \\
 = &\frac{(-1)^n \langle n \rangle_{n-1}}{(x-1)^{n+1}} \left(\ln x - \frac{x-1}{x} \right) + (-1)^{n-1} \sum_{k=1}^{n-1} \frac{(n-k)! \langle n \rangle_{k-1}}{x^{n-k+1}(x-1)^k} \\
 = &\frac{(-1)^n \langle n \rangle_n}{(x-1)^{n+1}} \ln x + (-1)^{n-1} \sum_{k=1}^n \frac{(n-k)! \langle n \rangle_{k-1}}{x^{n-k+1}(x-1)^k} \\
 = &(-1)^{n-1} n! \left[\sum_{k=1}^n \frac{1}{(n-k+1)x^{n-k+1}(x-1)^k} - \frac{1}{(x-1)^{n+1}} \ln x \right] \\
 = &(-1)^n \frac{n!}{x^{n+1}} \Phi \left(\frac{x-1}{x}, 1, n+1 \right).
 \end{aligned}$$

The formula (4) is thus proved.

From absolute monotonicity and Bernstein function property in Lemma 6, it is immediate to obtain that the function $f(x)$ is completely monotonic on $(0, \infty)$ and the functions $x^{n+1}f^{(n)}(x)$ for all $n \geq 0$ are Bernstein functions on $(0, \infty)$.

The inequality (5) follows from the complete monotonicity of $f(x)$ and the formula (3). The proof of Theorem 2 is complete. \square

Proof of Theorem 3. By (7), (16) in Lemma 7, and (4) in sequence, we obtain

$$\begin{aligned}
 F^{(m)}(t) &= [f(h_{1,-4}(t))]^{(m)} \\
 &= \sum_{\ell=0}^m f^{(\ell)}(x) B_{m,\ell}(h'_{1,-4}(t), h''_{1,-4}(t), \dots, h_{1,-4}^{(m-\ell+1)}(t)) \\
 &= 2^m \sum_{\ell=0}^m \frac{\ell!}{x^{\ell+1}} \Phi \left(\frac{x-1}{x}, 1, \ell+1 \right) [2(m-\ell)-1]!! \binom{2m-\ell-1}{2(m-\ell)} \frac{1}{(1-4t)^{m-\ell/2}} \\
 &= \frac{2^m}{(1-4t)^{m+1/2}} \sum_{\ell=0}^m \ell! [2(m-\ell)-1]!! \binom{2m-\ell-1}{\ell-1} \Phi \left(\frac{\sqrt{1-4t}-1}{\sqrt{1-4t}}, 1, \ell+1 \right).
 \end{aligned}$$

The absolute monotonicity of $F(t)$ on $(-\infty, \frac{1}{4})$ follows from the absolute monotonicity of the Lerch transcendent $\Phi(x, s, a)$ for $s, a > 0$ with respect to $x \in (-\infty, 1)$. \square

Proof of Theorem 4. It is well known that the Stirling numbers of the first kind $s(n, k)$ for $n \geq k \geq 1$ can be generated [4, 11, 13] by

$$\frac{[\ln(1+x)]^k}{k!} = \sum_{n=k}^{\infty} s(n, k) \frac{x^n}{n!}, \quad |x| < 1.$$

It can be rewritten as

$$\left[\frac{\ln(1+x)}{x} \right]^k = \sum_{n=0}^{\infty} \frac{s(n+k, k)}{\binom{n+k}{k}} \frac{x^n}{n!}, \quad |x| < 1.$$

This means that

$$\begin{aligned} \frac{s(n+k, k)}{\binom{n+k}{k}} &= \lim_{x \rightarrow 0} \frac{d^n}{d x^n} \left[\frac{\ln(1+x)}{x} \right]^k = \lim_{x \rightarrow 0} \frac{d^n}{d x^n} [f(x+1)]^k \\ &= \lim_{x \rightarrow 0} \sum_{\ell=0}^n \langle u^k \rangle_{\ell}^{(k)} B_{n, \ell}(f'(x+1), f''(x+1), \dots, f^{(n+\ell-1)}(x+1)) \\ &= \lim_{x \rightarrow 0} \sum_{\ell=0}^n \langle k \rangle_{\ell} u^{k-\ell} B_{n, \ell}(f'(x+1), f''(x+1), \dots, f^{(n+\ell-1)}(x+1)) \\ &= \sum_{\ell=0}^n \langle k \rangle_{\ell} f^{k-\ell}(1) B_{n, \ell}(f'(1), f''(1), \dots, f^{(n+\ell-1)}(1)) \\ &= \sum_{\ell=0}^n \langle k \rangle_{\ell} B_{n, \ell} \left(-\frac{1!}{2}, \frac{2!}{3}, \dots, (-1)^{n+\ell-1} \frac{(n+\ell-1)!}{n-\ell+2} \right) \\ &= (-1)^n \sum_{\ell=0}^n \langle k \rangle_{\ell} B_{n, \ell} \left(\frac{1!}{2}, \frac{2!}{3}, \dots, \frac{(n+\ell-1)!}{n-\ell+2} \right) \\ &= \sum_{\ell=0}^n (-1)^{\ell} \frac{\langle k \rangle_{\ell}}{\ell!} \sum_{m=1}^{\ell} (-1)^m \binom{\ell}{m} \frac{s(n+m, m)}{\binom{n+m}{m}}, \end{aligned}$$

where $u = f(x+1)$ and we used the identity (10) in the last step. The recurrence relation (6) is thus proved. \square

Remark 2. This paper is a slightly revised version of the preprint [20].

REFERENCES

1. N. Bourbaki, *Functions of a Real Variable, Elementary Theory*, Translated from the 1976 French original by Philip Spain. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2004; Available online at <https://doi.org/10.1007/978-3-642-59315-4>.

2. N. D. Cahill, J. R. D'Errico, D. A. Narayan, and J. Y. Narayan, *Fibonacci determinants*, *College Math. J.* **3** (2002), 221–225; Available online at <https://doi.org/10.2307/1559033>.
3. W. Cheney and W. Light, *A Course in Approximation Theory*, Reprint of the 2000 original, *Graduate Studies in Mathematics*, Volume **101**, American Mathematical Society, Providence, RI, 2009; Available online at <https://doi.org/10.1090/gsm/101>.
4. L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Revised and Enlarged Edition, D. Reidel Publishing Co., Dordrecht and Boston, 1974.
5. B.-N. Guo and F. Qi, *A property of logarithmically absolutely monotonic functions and the logarithmically complete monotonicity of a power-exponential function*, *Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys.* **72** (2010), no. 2, 21–30.
6. S. Hu and M.-S. Kim, *Two closed forms for the Apostol-Bernoulli polynomials*, arXiv preprint (2015), available online at <http://arxiv.org/abs/1509.04190>.
7. T. Kim and D. S. Kim, *Differential equations associated with Catalan–Daehee numbers and their applications*, *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM* **111** (2017), no. 4, 1071–1081; Available online at <https://doi.org/10.1007/s13398-016-0349-4>.
8. D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, 1993; Available online at <https://doi.org/10.1007/978-94-017-1043-5>.
9. F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark (eds.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York, 2010; Available online at <http://dlmf.nist.gov/>.
10. F. Qi, *Derivatives of tangent function and tangent numbers*, *Appl. Math. Comput.* **268** (2015), 844–858; Available online at <https://doi.org/10.1016/j.amc.2015.06.123>.
11. F. Qi, *Diagonal recurrence relations for the Stirling numbers of the first kind*, *Contrib. Discrete Math.* **11** (2016), 22–30; Available online at <http://hdl.handle.net/10515/sy5wh2dx6> and <https://doi.org/10515/sy5wh2dx6>.
12. F. Qi, *Diagonal recurrence relations, inequalities, and monotonicity related to the Stirling numbers of the second kind*, *Math. Inequal. Appl.* **19** (2016), no. 1, 313–323; Available online at <https://doi.org/10.7153/mia-19-23>.
13. F. Qi, *Integral representations and properties of Stirling numbers of the first kind*, *J. Number Theory* **133** (2013), no. 7, 2307–2319; Available online at <https://doi.org/10.1016/j.jnt.2012.12.015>.
14. F. Qi, V. Čerňanová, and Y. S. Semenov, *On tridiagonal determinants and the Cauchy product of central Delannoy numbers*, ResearchGate Working Paper (2016), available online at <https://doi.org/10.13140/RG.2.1.3772.6967>.
15. F. Qi and R. J. Chapman, *Two closed forms for the Bernoulli polynomials*, *J. Number Theory* **159** (2016), 89–100; Available online at <https://doi.org/10.1016/j.jnt.2015.07.021>.
16. F. Qi and B.-N. Guo, *A determinantal expression and a recurrence relation for the Euler polynomials*, *Adv. Appl. Math. Sci.* **16** (2017), no. 9, 297–309.

17. F. Qi and B.-N. Guo, *Explicit and recursive formulas, integral representations, and properties of the large Schröder numbers*, Kragujevac J. Math. **41** (2017), no. 1, 121–141.
18. F. Qi and B.-N. Guo, *Expressing the generalized Fibonacci polynomials in terms of a tridiagonal determinant*, Matematiche (Catania) **72** (2017), no. 1, 167–175; Available online at <https://doi.org/10.4418/2017.72.1.13>.
19. F. Qi and B.-N. Guo, *Some determinantal expressions and recurrence relations of the Bernoulli polynomials*, Mathematics **4** (2016), no. 4, Article 65, 11 pages; Available online at <https://doi.org/10.3390/math4040065>.
20. F. Qi and B.-N. Guo, *Some properties of a solution to a family of inhomogeneous linear ordinary differential equations*, Preprints **2016**, 2016110146, 11 pages; Available online at <https://doi.org/10.20944/preprints201611.0146.v1>.
21. F. Qi and B.-N. Guo, *Two nice determinantal expressions and a recurrence relation for the Apostol–Bernoulli polynomials*, J. Indones. Math. Soc. **23** (2017), no. 1, 81–87; Available online at <https://doi.org/10.22342/jims.23.1.274.81-87>.
22. F. Qi and W.-H. Li, *Integral representations and properties of some functions involving the logarithmic function*, Filomat **30** (2016), no. 7, 1659–1674; Available online at <https://doi.org/10.2298/FIL1607659Q>.
23. F. Qi, M. Mahmoud, X.-T. Shi, and F.-F. Liu, *Some properties of the Catalan–Qi function related to the Catalan numbers*, SpringerPlus (2016), **5**:1126, 20 pages; Available online at <https://doi.org/10.1186/s40064-016-2793-1>.
24. F. Qi, X.-T. Shi, and B.-N. Guo, *Two explicit formulas of the Schröder numbers*, Integers **16** (2016), Paper No. A23, 15 pages.
25. F. Qi, X.-T. Shi, F.-F. Liu, and D. V. Kruchinin, *Several formulas for special values of the Bell polynomials of the second kind and applications*, J. Appl. Anal. Comput. **7** (2017), no. 3, 857–871; Available online at <https://doi.org/10.11948/2017054>.
26. F. Qi, J.-L. Wang, and B.-N. Guo, *A recovery of two determinantal representations for derangement numbers*, Cogent Math. (2016), **3**: 1232878, 7 pages; Available online at <https://doi.org/10.1080/23311835.2016.1232878>.
27. F. Qi, J.-L. Zhao, and B.-N. Guo, *Closed forms for derangement numbers in terms of the Hessenberg determinants*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **112** (2018), in press; Available online at <http://dx.doi.org/10.1007/s13398-017-0401-z>.
28. R. L. Schilling, R. Song, and Z. Vondraček, *Bernstein Functions—Theory and Applications*, 2nd ed., de Gruyter Studies in Mathematics **37**, Walter de Gruyter, Berlin, Germany, 2012; Available online at <https://doi.org/10.1515/9783110269338>.
29. C.-F. Wei and F. Qi, *Several closed expressions for the Euler numbers*, J. Inequal. Appl. **2015**, **2015**:219, 8 pages; Available online at <https://doi.org/10.1186/s13660-015-0738-9>.
30. D. V. Widder, *The Laplace Transform*, Princeton University Press, Princeton, 1946.
31. A.-M. Xu and G.-D. Cen, *Closed formulas for computing higher-order derivatives of functions involving exponential functions*, Appl. Math. Comput. **270** (2015), 136–141; Available online at <http://dx.doi.org/10.1016/j.amc.2015.08.051>.

32. A.-M. Xu and Z.-D. Cen, *Some identities involving exponential functions and Stirling numbers and applications*, J. Comput. Appl. Math. **260** (2014), 201–207; Available online at <http://dx.doi.org/10.1016/j.cam.2013.09.077>.

Feng Qi

Institute of Mathematics
Henan Polytechnic University
Jiaozuo, Henan, 454010
China;
College of Mathematics
Inner Mongolia University for Nationalities
Tongliao, Inner Mongolia, 028043
China
Department of Mathematics
College of Science
Tianjin Polytechnic University
Tianjin, 300387
China
E-mail: qifeng618@gmail.com
qifeng618@hotmail.com
qifeng618@qq.com
URL: <https://qifeng618.wordpress.com>

(Received 05.04.2017)

(Revised 01.10.2017)

Bai-Ni Guo

School of Mathematics and Informatics
Henan Polytechnic University
Jiaozuo City, 454010
Henan Province
China
E-mail: bai.ni.guo@gmail.com
bai.ni.guo@hotmail.com