

CHROMATIC ZEROS AND THE GOLDEN RATIO

Saeid Alikhani, Yee-hock Peng

In this note, we investigate τ^n , where $\tau = \frac{1 + \sqrt{5}}{2}$ is the golden ratio as chromatic roots. Using some properties of FIBONACCI numbers, we prove that $\tau^n (n \in \mathbb{N})$, cannot be roots of any chromatic polynomial.

1. INTRODUCTION

Let $G = (V, E)$ be a simple graph and $\lambda \in \mathbb{N}$. A mapping $f: V \rightarrow \{1, 2, \dots, \lambda\}$ is called a λ -colouring of G if $f(u) \neq f(v)$ whenever the vertices u and v are adjacent in G . The number of distinct λ -colourings of G , denoted by $P(G, \lambda)$ is called the *chromatic polynomial* of G . A root of $P(G, \lambda)$ is called a *chromatic root* of G . An interval is called a *root-free interval* for a chromatic polynomial $P(G, \lambda)$ if G has no chromatic root in this interval. It is well-known that $(-\infty, 0)$ and $(0, 1)$ are two maximal root-free intervals for the family of all graphs (see [2]). JACKSON [2] showed that $(1, 32/27]$ is another maximal root-free interval for the family of all graphs and the value $32/27$ is best possible.

A *plane graph* is a graph drawn in the plane in such a way that any pair of edges meet only once at their end vertices. A *plane triangulation* is a loopless plane graph in which all faces have size three. A *near-triangulation* graph, is a plane graph with at most one non-triangular face.

We recall that a complex number ζ is called an *algebraic number* (resp., an *algebraic integer*) if it is a root of some monic polynomial with rational (resp., integer) coefficients (see [6]). Since the chromatic polynomial $P(G, \lambda)$ is a monic polynomial in λ with integer coefficients, its roots are, by definition, algebraic integers. This naturally raises the question: *Which* algebraic integers can occur as zeros of chromatic polynomials? Clearly those lying in $(-\infty, 0) \cup (0, 1) \cup (1, 32/27]$ are forbidden.

Using this reasoning, TUTTE [7] proved that $B_5 = \frac{3 + \sqrt{5}}{2} = 1 + \tau = \tau^2$ ($B_n = 2 + 2 \cos(2\pi/n)$ is the n -th *Beraha number* [1]) cannot be a chromatic root,

2000 Mathematics Subject Classification. 05C15, 11B39.

Keywords and Phrases. Chromatic zero, FIBONACCI numbers, golden ratio.

for otherwise $B_5^* = \frac{3 - \sqrt{5}}{2}$ would also be a chromatic root, which is impossible, since $B_5^* \in (0, 1)$. SALAS and SOKAL in [5] extended this result to show that the generalized BERAHA numbers $B_n^{(k)} = 4 \cos^2(k\pi/n)$ for $n = 5, 7, 8, 9$ and $n \geq 11$, with k coprime to n , are never chromatic roots. For $n = 10$ they showed the weaker result that $B_{10} = \frac{5 + \sqrt{5}}{2}$ and $B_{10}^* = \frac{5 - \sqrt{5}}{2}$ are not chromatic roots of any plane near-triangulation. The results and conjectures in [3, 8, 9, 10] show that τ and τ^2 has a special significance for chromatic polynomials of plane triangulations.

In this note we investigate τ^n as chromatic zeros. We use some properties of FIBONACCI numbers and the golden ratio to prove that $\tau^n (n \in \mathbb{N})$, where $\tau = \frac{1 + \sqrt{5}}{2}$ is the golden ratio, cannot be roots of any chromatic polynomials.

2. FIBONACCI NUMBERS AND THE GOLDEN RATIO

FIBONACCI numbers are terms of the sequence defined in a quite simple recursive fashion. However, despite their simplicity, they have some curious properties which are worth attention. In this section, we look at some of the important features of these numbers which are needed to prove the main result in next section.

Definition 1. *The sequence F_n of natural numbers defined by the equations $F_0 = 0$, $F_1 = 1$ and $F_n = F_{n-1} + F_{n-2}$ ($n \geq 2$) is called the Fibonacci sequence or the sequence of the Fibonacci numbers. The n -th term in the sequence is called the n -th Fibonacci number.*

We recall the following theorem:

Theorem 1 ([4], p. 78) $\tau^n = F_n \tau + F_{n-1}$ ($n \in \mathbb{N}$).

Corollary 1. *For every $n \in \mathbb{N}$, $|\tau^{-1} F_n + F_{n-1}| < 1$.*

Proof. Similar to theorem 1, by induction on n , we have $(1 - \tau)F_n + F_{n-1} = (1 - \tau)^n$. Therefore, we have the result.

3. CHROMATIC ZEROS AND THE GOLDEN RATIO

In this section, we prove that τ^n can not be a chromatic root for any natural number n . We state and prove following theorem:

Theorem 2. *Suppose that a, b are rational numbers, and $r \geq 2$ is an integer that is not a perfect square, and $a - |b|\sqrt{r} < 32/27$. Then $a + b\sqrt{r}$ is not the root of any chromatic polynomial.*

Proof. If $\lambda = a + b\sqrt{r}$ is a root of some polynomial with integer coefficients (e.g. a chromatic polynomial), then so is $\lambda^* = a - b\sqrt{r}$. But λ or λ^* can not belong to $(-\infty, 0) \cup (0, 1) \cup (1, 32/27]$, a contradiction. \square

The following theorem is our main result:

Theorem 3. $\tau^n (n \in \mathbb{N})$ is not a root of any chromatic polynomials.

Proof. By Theorem 1, we can consider τ^n of the form $\lambda = a + b\sqrt{r}$ with $a = \frac{F_n}{2} + F_{n-1}$, $b = \frac{F_n}{2}$, and $r = 5$. Since $a - |b|\sqrt{r} = -\tau^{-1}F_n + F_{n-1}$, by Corollary 1, $a - |b|\sqrt{r} < 1 < 32/27$. Therefore we have the result by Theorem 2. \square

Acknowledgment. The authors wish to thank the referee who made suggestions, and shortened our manuscript.

REFERENCES

1. S. BERAHA: *Infinite non-trivial families of maps and chromials*. Ph.D. thesis, John Hopkins University, 1975.
2. B. JACKSON: *A zero free interval for chromatic polynomials of graphs*. *Combin. Probab. Comput.*, **2** (1993), 325–336.
3. B. JACKSON: *Zeros of chromatic and flow polynomials of graphs*. *Journal of Geometry*, **76**, No. 1–2 (2003), 95–109.
4. T. KOSHY: *Fibonacci and Lucas numbers with applications*. A Willey-Interscience Publication, 2001.
5. J. SALAS, A. D. SOKAL: *Transfer matrices and partition function zeros for anti-ferromagnetic Potts models*. I. General Theory and squar lattice polynomial. *J. Statist. Phys.*, **104** (2001), 609–699.
6. I. STEWART, D. TALL: *Algebraic Number Theory*, 2nd ed, Chapman and Hall, London-New York, 1987.
7. W. T. TUTTE: *On chromatic polynomials and the golden ratio*. *J. Combinatorial Theory*, Ser B **9** (1970), 289–296.
8. W. T. TUTTE: *The graph of the chromial of a graph*. *Combinatorial Mathematics III*, Lecture Notes in Mathematics, Volume **452** (1975) 55-61, Springer, 1975
<http://www.springerlink.com/content/h834255pq053trj>
9. W. T. TUTTE: *The golden ratio in the theory of chromatic polynomials*. *Annals of the New York Academy of Sciences*, **175**, No. 1, (1970), 391–402.
10. W. T. TUTTE: *More about chromatic polynomials and the golden ratio*. *Combinatorial Structures and their Applications* (R.K.Guy, ed.), Proc. Calgary Internat. Conf., Calgary, 1969, Gordon and Breach, P. 439–453.

Department of Mathematics, Faculty of Science,
Yazd University,
89195-741, Yazd, Iran

Institute for Mathematical Research,
University Putra Malaysia,
43400 UPM Serdang, Malaysia
E-mail: alikhani206@gmail.com

(Received May 3, 2008)
(Revised August 26, 2008)

Institute for Mathematical Research and Department of Mathematics,
University Putra Malaysia,
43400 UPM Serdang, Malaysia
E-mail: yhpeng@science.upm.edu.my